

NuLytely/GoLytely and Magnesium Citrate 2-Day Colonoscopy Prep



 **PRESBYTERIAN**

IMPORTANT!
**Please read these
instructions one week
prior to your procedure**

It will take you several days to prepare for this test. You may also need to stop taking certain medications up to a week in advance. Please read these instructions well in advance of your procedure date. We recommend that you post these instructions in a visible location at home to remind you of the various steps you will need to take.

- **You will need to take the day off from work.**
- **You will need someone to drive you home after your procedure. You cannot drive yourself home because you will be receiving sedation medication.**
- **If your doctor removes polyps during the procedure, we recommend that you do not travel for two weeks after the procedure.**

Presbyterian Medical Group
Gastroenterology
(505) 224-7000

Patient responsibility

- If there are any changes with your insurance(s), please notify our office prior to your procedure date.
- After your procedure is scheduled, you may receive a call from the Presbyterian Registration department to verify your information and insurance coverage. You may call the department with registration questions at **559-1009**.
- If you need financial assistance with the cost of the procedure, or if you have financial or billing questions, please call 923-6600.
- If your insurance requires a referral, it must be current at the time of your procedure, or you may be liable for payment.
- Remember to bring your insurance card(s) with you the day of your procedure.

Purchases

- One 10-oz. bottle of Magnesium Citrate. This is available over the counter; you do not need a prescription.
- Fill the prescription for NuLytely/GoLytely you were given. You may purchase plain or pineapple flavored NuLytely/GoLytely.

Five days prior to your colonoscopy

- **Coumadin, Plavix, Persantine, Ticlid (Blood Thinners):**

Do not discontinue these medications without instructions from your cardiologist, prescribing physician or Coumadin clinic. Contact the prescribing physician for instructions. If your physician does not want you to stop these medications, please notify our office at 224-7000.

- If you are diabetic, please contact your prescribing provider for diabetic medication instructions.

DO NOT take the following medications:

- **Alka Seltzer, Pepto Bismol, Iron supplements**

Three days prior to your colonoscopy

DO NOT eat foods that contain seeds, corn, or nuts

Two days prior to your colonoscopy

- **DO NOT EAT ANY SOLID FOODS!**

- You can only drink clear liquids for breakfast, lunch, and dinner. See the list of clear liquids in this packet. Avoid red, orange and purple colored Jell-O and Gatorade. This discolors your stool and interferes with the colonoscopy.

- **At 5:30 p.m. drink the entire 10-oz. bottle of magnesium citrate.**

The day before your colonoscopy

- **DO NOT EAT ANY SOLID FOODS!**

- Continue drinking clear liquids only throughout the day and evening.
- We suggest that you mix the NuLytely/GoLytely in the morning and put it in the refrigerator. It tends to taste better when it is cold.

The night before

- At 6 p.m. begin drinking the NuLytely/GoLytely at a rate of one glass every 15 minutes until you have taken half (2 liters) of the solution.
If you become nauseated or begin vomiting while drinking the solution, slow down. When the nausea subsides, begin drinking the solution again.

On the day of your colonoscopy

- 6 hours before your procedure time, drink the remaining NuLytely/GoLytely solution (2 liters)
- By 6 a.m. the day of your procedure, please take your blood pressure, cardiac and seizure medications.
- You may have clear liquids up until 4 hours prior to your appointment time.
- Bring inhalers, reading glasses, and hearing aids if you use them.
- Remove all jewelry and leave valuables at home.

Normal symptoms you may experience from the preparation:

- You will have diarrhea from the bowel preparation medications. This will start between a few minutes to three hours after you start the prep.
- Because of the diarrhea, you will need access to a toilet once you begin the bowel preparation. Plan your day with this need in mind.
- Most people have bloating and abdominal discomfort. This is normal. Do not be alarmed if you feel these symptoms.
- If you become nauseated or begin vomiting while drinking the solution, slow down. When the nausea subsides, begin drinking the solution again. Some people do not like the taste or the smell of the medicine, try chilling the liquids. Please take the medicine as directed.

Clear Liquid Diet

You may only drink or eat items on this list.

- Water
- Tea (no milk or cream; sugar is permitted.)
- Carbonated beverages (soft drinks – regular or sugar free.)
- Apple juice
- White grape juice
- Gatorade (no red, purple or orange in color)
- Lemonade (no pulp)
- Limeade (no pulp)
- Clear chicken or beef broth (no meat, vegetables, noodles, rice)
- Bouillon
- Clear consommé
- Snowcones or popsicles (no red, purple or orange in color)
- Hard candy
- Jell-O (no red, purple, or orange in color. We prefer that you have only light colors such as yellow or pink.)
- **ALCOHOL IS NOT PERMITTED!**

Call our office at
224-7000 if:

- **You cannot complete the bowel preparation as ordered**
- **You have the following symptoms after your procedure:**
 - Nausea
 - Vomiting
 - Fever
 - Chills
 - Bleeding
 - Abdominal pain

We require that you contact us at least 72 hours before your scheduled test if you wish to cancel your test.



Presbyterian Medical Group
Gastroenterology

(505) 224-7000