

**Presbyterian Senior Care (HMO)/(HMO-POS)
Presbyterian UltraFlex (HMO-POS)
Presbyterian MediCare PPO
Presbyterian Dual Plus (HMO D-SNP)
Formulary Prior Authorization Criteria
Effective July 1, 2024**

The formulary may change at any time. You will receive notice when required.

For the most recent list of drugs, information on obtaining a coverage determination or exception, or other questions, please contact the Presbyterian Customer Service Center.

Presbyterian Senior Care, Presbyterian UltraFlex, and Presbyterian MediCare PPO:



(505) 923-6060
1-800-797-5343
(TTY 711)



October 1 to March 31:

8 a.m. to 8 p.m., seven days a week
(except holidays)

April 1 to September 30:

8 a.m. to 8 p.m., Monday - Friday
(except holidays)

Presbyterian Dual Plus:



(505) 923-7675
1-855-465-7737
(TTY 711)



www.phs.org/Medicare

Learn more about Presbyterian's Nondiscrimination Notice and Interpreter Services.

Based on a Model of Care review, Presbyterian Dual Plus (HMO D-SNP) has been approved by the National Committee for Quality Assurance (NCQA) to operate a Special Needs Plan (SNP) through 2025.

Abilify MyCite (aripiprazole with sensor)

Products Affected

- Abilify MyCite
- Abilify MyCite Maintenance Kit Oral Tablet Therapy Pack 15 MG, 2 MG, 20 MG, 5 MG
- Abilify MyCite Starter Kit Oral Tablet Therapy Pack 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	Patients with dementia-related psychosis.
Required Medical Information	Chart notes documenting that the patient has tried at least two (2) oral atypical anti-psychotics, one of which must be aripiprazole.
Age Restrictions	Patient is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Initial: 3 months Renewal: 6 months
Other Criteria	Reauthorization: Documentation that the patient is clinically stable on Abilify MyCite and the prescriber documents that the patient requires the continued use of Abilify MyCite.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Abilify ODT (aripiprazole)

Products Affected

- ARIPiprazole Oral Tablet Dispersible

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation that the patient has tried/failed aripiprazole tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Actemra (tocilizumab SubQ)

Products Affected

- Actemra ACTPen
- Actemra Subcutaneous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Rheumatoid Arthritis (RA): Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. An adequate trial (3 months or more) of one of the following other DMARDs: hydroxychloroquine, leflunomide, methotrexate, sulfasalazine. Inadequate response to two of the following, unless contraindicated or not tolerated: Amjevita, Enbrel, Hadlima, Rinvoq, Xeljanz. 2. Adult patients with Giant Cell Arteritis (GCA): trial and failure, unless contraindicated or not tolerated, of oral corticosteroids.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	Continuation of therapy: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	Yes

Actiq (fentanyl transmucosal)

Products Affected

- fentaNYL Citrate Buccal Lozenge On A Handle 1200 MCG, 400 MCG, 600 MCG, 800 MCG
- FentaNYL Citrate Buccal Lozenge On A Handle 1600 MCG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Trial and failure of an immediate release oral opiate and must be used in combination with a long-acting oral opiate.
Age Restrictions	16 years or older
Prescriber Restrictions	Oncologist or Pain Specialist
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adcirca (tadalafil)

Products Affected

- Alyq
- Tadalafil (PAH)

PA Criteria	Criteria Details
Exclusion Criteria	Tadalafil is excluded from coverage for the treatment of Erectile Dysfunction.
Required Medical Information	Documentation of Pulmonary Arterial Hypertension as determined by a right heart catheterization.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adempas (riociguat)

Products Affected

- Adempas

PA Criteria	Criteria Details
Exclusion Criteria	Must not be used in combination with nitrates, nitric oxide donors, or PDE-5 inhibitors.
Required Medical Information	If using for pulmonary arterial hypertension (PAH), must have tried and failed or have a contraindication to Revatio (sildenafil) or Adcirca (tadalafil).
Age Restrictions	18 years or older
Prescriber Restrictions	Cardiologist or Pulmonologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Afinitor (everolimus)

Products Affected

- Everolimus Oral Tablet 10 MG, 2.5 MG, 5 MG, 7.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Afinitor Disperz (everolimus)

Products Affected

- Everolimus Oral Tablet Soluble

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Akeega (niraparib and abiraterone)

Products Affected

- Akeega

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Alecensa (alectinib)

Products Affected

- Alecensa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Alunbrig (brigantinib)

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Amjevita (adalimumab-atto)

Products Affected

- Amjevita Subcutaneous Solution Auto-Injector
- Amjevita Subcutaneous Solution Prefilled Syringe 40 MG/0.4ML
- Amjevita-Ped 10kg to <15kg
- Amjevita-Ped 15kg to <30kg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Ankylosing Spondylitis (AS): peripheral arthritis must have a trial of sulfasalazine and an NSAID. Patients with axial disease and failure of NSAIDs can be started without a trial of sulfasalazine. 2. Crohn's disease (CD): Inadequate response or intolerance to conventional therapy. For the purpose of this policy, conventional therapy includes the use of ONE of the following: i. Corticosteroids (e.g., prednisone, prednisolone, dexamethasone, budesonide). ii. Methotrexate iii. Thiopurines (azathioprine, mercaptopurine). 3. Juvenile Idiopathic Arthritis (JIA): An adequate trial (3 months or more) of one of the following DMARDs: i. Leflunomide ii. Methotrexate iii. Sulfasalazine. 4. Plaque Psoriasis (PsO): a. The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to, or is intolerant, a 3-month trial of a topical agent (topical corticosteroid, topical calcineurin inhibitor, topical vitamin D analogs, etc.). 5. Psoriatic Arthritis (PsA): at least a three month trial of one of the following: cyclosporine, leflunomide (LEF), MTX, SSZ 6. Rheumatoid Arthritis (RA): Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. c. An adequate trial (3 months or more) of one of the following other DMARDs: i. Hydroxychloroquine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine 7. Ulcerative Colitis (UC): The member has had an inadequate response to one of the following: aminosalicylates, corticosteroids, thiopurines, or cyclosporine.</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist, dermatologist, gastroenterologist, or ophthalmologist.
Coverage Duration	One (1) year
Other Criteria	8. Hidradenitis Suppurativa (HS): Documented diagnosis of Hurley Stage III HS or refractory Hurley Stage II hidradenitis suppurativa. 9. Uveitis: a. Documented diagnosis of non-infectious intermediate, posterior and panuveitis in adult patients and meets the following: i. documented trial failure, contraindication, or intolerance to conventional therapy such as ophthalmic or systemic corticosteroids AND immunosuppressive drugs (e.g., azathioprine, cyclosporine, methotrexate, or tacrolimus). For all diagnoses: 1. Current PPD (tuberculosis) negative skin test or negative QuantiFERON-TB Gold test before the initiation of therapy, 2. The appropriate Disease Specific Criteria has been met. CONTINUATION CRITERIA: Documentation of positive response with treatment
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Androderm (testosterone topical)

Products Affected

- Androderm Transdermal Patch 24 Hour

PA Criteria	Criteria Details
Exclusion Criteria	Testosterone replacement will not be covered for the treatment of sexual dysfunction.
Required Medical Information	Trial and failure of Androgel (testosterone gel). If using for primary hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with elevated luteinizing hormone (LH) and follicular stimulating hormone (FSH) levels. If using for hypogonadotropic hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with low to low-normal LH and FSH levels.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AndroGel 1.62% (testosterone topical)

Products Affected

- Testosterone Transdermal Gel 1.62 %, 10 MG/ACT (1.62%), 25 MG/2.5GM (1%), 12.5 MG/ACT (2%), 12.5 MG/ACT (1%), 40.5 MG/2.5GM (1.62%), 50 MG/5GM (1%), 20.25 MG/1.25GM (1.62%), 20.25 (1%)

PA Criteria	Criteria Details
Exclusion Criteria	Testosterone replacement will not be covered for the treatment of sexual dysfunction.
Required Medical Information	If using for primary hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with elevated luteinizing hormone (LH) and follicular stimulating hormone (FSH) levels. If using for hypogonadotropic hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with low to low-normal LH and FSH levels.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Apokyn (apomorphine)

Products Affected

- Apokyn Subcutaneous Solution Cartridge
- Apomorphine HCl Subcutaneous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Must be used as an adjunct to levodopa and one (1) formulary oral dopamine agonist medication indicated for Parkinson's disease.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Arcalyst (rilonacept)

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Exclusion Criteria	Will not be used in combination with etanercept, adalimumab, anakinra, abatacept, or infliximab.
Required Medical Information	
Age Restrictions	12 years or older for the indications of Cryopyrin-Associated Periodic Syndromes, including Familial Cold Autoinflammatory Syndrome, and Muckle-Wells Syndrome, recurrent pericarditis. Adults and pediatrics weighing 10 kg or more for the maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist.
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Aubagio (teriflunomide)

Products Affected

- Teriflunomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Augtyro (repotrecitinib)

Products Affected

- Augtyro

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Austedo (deutetrabenazine)

Products Affected

- Austedo
- Austedo XR
- Austedo Patient Titration Kit
- Austedo XR Patient Titration

PA Criteria	Criteria Details
Exclusion Criteria	Deutetrabenazine is not covered for patients who are actively suicidal, who have untreated or inadequately treated depression, who have impaired hepatic function, or who are currently taking monoamine oxidase inhibitors or reserpine.
Required Medical Information	Documentation that member is being monitored for depression and suicidal ideation. Chorea associated with Huntington disease (HD): Documentation that the patient is ambulatory and a trial and failure of tetrabenazine. Tardive Dyskinesia (TD): Documentation of a baseline Abnormal Involuntary Movement Scale (AIMS) must be provided.
Age Restrictions	
Prescriber Restrictions	HD: Neurologist TD: Psychiatrist or Neurologist
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	Renewal: Chart notes documenting that the patient's disease has improved based on prescriber's assessment while on therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Auvelity (dextromethorphan/bupropion)

Products Affected

- Auvelity

PA Criteria	Criteria Details
Exclusion Criteria	1. Seizure disorder. 2. Current or prior diagnosis of bulimia or anorexia nervosa. 3. Abrupt discontinuation of alcohol, benzodiazepines, barbiturates, and antiepileptic drugs. 4. Use with an MAOI.
Required Medical Information	1. Member has a clinical diagnosis of major depressive disorder (MDD) as defined by DSM-5 criteria and/or appropriate depression rating scale (e.g., MADRS). 2. Member has had previous treatment, contraindication, or intolerance to at least two antidepressants from two different classes (e.g., SSRI, SNRI).
Age Restrictions	18 years of age and over
Prescriber Restrictions	
Coverage Duration	Initial: 6 months. Continuation: 1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Ayvakit (avapritinib)

Products Affected

- Ayvakit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Balversa (erdafitinib)

Products Affected

- Balversa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Benlysta (belimumab)

Products Affected

- Benlysta Subcutaneous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Must be used in combination with standard therapy.
Age Restrictions	
Prescriber Restrictions	Nephrologist or Rheumatologist
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	Documentation must be submitted demonstrating a clinical benefit has been established and maintained compared to baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Besremi

Products Affected

- Besremi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Bosulif (bosutinib)

Products Affected

- Bosulif

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Braftovi (encorafenib)

Products Affected

- Braftovi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Bronchitol (mannitol)

Products Affected

- Bronchitol

PA Criteria	Criteria Details
Exclusion Criteria	Failure to pass Bronchitol tolerate test (BTT)
Required Medical Information	1. Diagnosis of cystic fibrosis. 2. Documentation of in adequate response to hypertonic saline and Pulmozyme, unless contraindicated or not tolerated. 3. Documentation that member has successfully complete the Bronchitol tolerance test (BTT).
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist.
Coverage Duration	One (1) year
Other Criteria	Continuation criteria: positive clinical response to therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Brukinsa (Zanubrutinib)

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Buphenyl (sodium phenylbutyrate)

Products Affected

- Sodium Phenylbutyrate Oral Powder 3 GM/TSP
- Sodium Phenylbutyrate Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cabometyx (cabozantinib)

Products Affected

- Cabometyx

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Calquence (acalabrutinib)

Products Affected

- Calquence

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Caplyta (lumateperone)

Products Affected

- Caplyta

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Schizophrenia (must meet all) - A. Diagnoses of schizophrenia. B. Member meets one of the following (i or ii: i. Failure of two of the following generic atypical antipsychotics at up to maximally tolerated doses, each used for at least 4 weeks, unless clinically significant adverse effects experienced or are contraindicated: aripiprazole, olanzapine, quetiapine, risperidone, ziprasidone, or, ii. member has diabetes mellitus for body mass index (BMI) greater than 30. 2. Bipolar Disorder (must meet all) - A. Diagnosis of bipolar disorder. B. Failure of two preferred atypical antipsychotics at up to maximally indicated doses used for at least 4 weeks, unless contraindicated or not tolerated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	Continuation criteria: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Caprelsa (vandetanib)

Products Affected

- Caprelsa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Carbaglu (carglumic acid)

Products Affected

- Carglumic Acid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cayston (aztreonam lysine)

Products Affected

- Cayston

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FEV1 must be 25% to 75% of predicted. Patient must have a positive sputum culture for Pseudomonas aeruginosa.
Age Restrictions	7 years or older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CGRP Inhibitor

Products Affected

- Aimovig

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Episodic Migraines or Chronic Migraines. History of failure or intolerance to two (2) preventive migraine medications from at least two (2) of the following classes: anti-depressant (e.g. venlafaxine), anti-convulsant (e.g. topiramate, divalproex) or anti-hypertensive (e.g. propranolol, verapamil).
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with a Headache Specialist, a Neurologist, or a Pain Specialist.
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cinryze (C1 inhibitor-human)

Products Affected

- Cinryze

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Clovique (trientine)

Products Affected

- Trientine HCl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented intolerance to penicillamine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Colony Stimulating Factor

Products Affected

- Udenyca Subcutaneous Solution Prefilled Syringe
- Zarxio

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Indications for approval: 1. Cancer patients receiving myelosuppressive therapy. 2. Patients with acute myeloid leukemia receiving induction or consolidation chemotherapy. 3. Cancer patients receiving bone marrow transplant. 4. Patients undergoing peripheral blood progenitor cell collection and therapy. 5. Patients with severe chronic neutropenia (cyclic or idiopathic) that meets one of the following: a) Documented diagnosis of severe chronic neutropenia (idiopathic) with an ANC of less than 500/mm ³ on three separate occasions over the previous 6 months. OR b) Documented diagnosis of severe chronic neutropenia (cyclic) with five consecutive days per cycle with an ANC less than 500/mm ³ for each of 3 regularly spaced cycles over a 6-month period. 6. Patients with severe chronic neutropenia (congenital) that have a documented diagnosis of congenital neutropenia. 7. For the treatment of hematopoietic syndrome of acute radiation symptoms.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Cometriq (cabozantinib)

Products Affected

- Cometriq (100 MG Daily Dose)
- Cometriq (140 MG Daily Dose)
- Cometriq (60 MG Daily Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Copiktra (duvelisib)

Products Affected

- Copiktra

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Corlanor (ivabradine)

Products Affected

- Corlanor

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Adults: Documented diagnosis of symptomatic chronic heart failure with left ventricular ejection fraction (LVEF) 35% or less, who are in normal sinus rhythm with a resting heart rate of at least 70 beats per minute (documented in the last 60 days) and are on a beta-blocker or have a contraindication to beta-blocker use. Documentation that the patient has tried/failed sacubitril/valsartan. Pediatrics: Documented diagnosis of Symptomatic Heart Failure due to Dilated Cardiomyopathy in patients 6 months of age or older and the patient is in normal sinus rhythm with an elevated heart rate and LVEF 45% or less. Documentation of a resting heart rate of 70-105 beats per minute.
Age Restrictions	
Prescriber Restrictions	Cardiologist or in consultation with a Cardiologist or a cardiac care specialist
Coverage Duration	1 year
Other Criteria	Continuation: Documentation of successful response to the medication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cosentyx (secukinumab)

Products Affected

- Cosentyx (300 MG Dose)
- Cosentyx Sensoready (300 MG)
- Cosentyx Sensoready Pen
- Cosentyx Subcutaneous
- Cosentyx UnoReady

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Plaque Psoriasis (PsO): a. The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to, or is intolerant to a 3-month trial of a topical agent (topical corticosteroid, topical calcineurin inhibitor, topical vitamin D analogs, etc.). 2. Psoriatic Arthritis (PsA): a. An adequate trial (3 months or more) of one of the following DMARDs: i. Cyclosporine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine. 3. Ankylosing Spondylitis (AS) - Trial and failure of NSAID. Patient s with peripheral arthritis must have a trial of sulfasalazine. Patients with axial disease and failure of NSAIDs can be started without a trial of sulfasalazine. 4. Non-radiographic Axial Spondyloarthritis: The patient has had a documented trial and failure of a non-steroidal anti-inflammatory drug (NSAID) or such treatment is contraindicated or not tolerated</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Current PPD (tuberculosis) negative skin test or negative Quantiferon-TB gold test prior to initiation of therapy. Continuation: Documentation of positive clinical response to therapy.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cotellic (cobimetinib)

Products Affected

- Cotellic

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Cystaran (cysteamine) ophthalmic solution

Products Affected

- Cystaran

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Original diagnosis made by an eye specialist. Individual has medical record documentation of a confirmed diagnosis of cystinosis. Individual has medical record documentation of corneal cystine crystals.
Age Restrictions	
Prescriber Restrictions	Ophthalmologist
Coverage Duration	6 months
Other Criteria	Reauthorization: documentation of positive clinical response to Cystaran therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daliresp (roflumimast)

Products Affected

- Roflumilast

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Patient must be 18 years of age or older. 2. Patient must have a diagnosis of severe COPD with chronic bronchitis (GOLD Stage III or worse). 3. Severe COPD is defined by the GOLD guidelines as FEV1 less than 50% predicted. 4. Patient must be currently receiving two standard treatments for severe COPD (i.e. long-acting beta-agonist, long-acting anticholinergic, and short-acting anticholinergic).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daraprim (pyrimethamine)

Products Affected

- Pyrimethamine Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Toxoplasmosis: documentation that the patient will be using a sulfonamide. Toxoplasmosis prophylaxis.
Age Restrictions	
Prescriber Restrictions	Infectious Disease or in consultation with Infectious Disease
Coverage Duration	8 weeks
Other Criteria	Pyrimethamine is not recommended for the treatment of acute malarial attacks and is not included in the CDC recommendations for the treatment of malaria. Pyrimethamine is not the drug of choice for malaria prophylaxis.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Daurismo (glasdegib)

Products Affected

- Daurismo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Deferasirox (Jadenu and Exjade)

Products Affected

- Deferasirox Oral Tablet
- Deferasirox Oral Tablet Soluble

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than 40 ml/min/1.73, patients with poor performance status, patients with high-risk myelodysplastic syndrome (MDS), and patients with advanced malignancies.
Required Medical Information	1. Chronic iron overload due to blood transfusions: The member has received a transfusion of at least 100 mL/kg of packed red blood cells (e.g., at least 20 units of packed red blood cells for a 40 kg person or more than 20 units in an individual weighing more than 40 kg), and serum ferritin is consistently greater than 1,000 mcg/L. 2. Iron overload in non-transfusion-dependent thalassemia syndromes (NTDT): The member has a liver iron concentration (LIC) of at least 5 mg Fe/g of dry weight and a serum ferritin greater than 300 mcg/L. Baseline lab values required for both indications: renal function, serum transaminases and bilirubin, and auditory and ophthalmic examinations.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months. Continuation: 1 year
Other Criteria	Continuation criteria: 1. Chronic iron overload due to blood transfusions: routine (monthly) blood counts, liver function, renal function, and ferritin. 2. NTDT: LIC every 6 months, routine (monthly) blood counts, liver function, renal function, and ferritin.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Depen (penicillamine tablets)

Products Affected

- penicillAMINE Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystinuria: Documented trial and failure or intolerance to Thiola (tiopronin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	Renewal: Documentation must be submitted demonstrating a successful response to the medication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dificid (fidaxomicin)

Products Affected

- Dificid Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Trial and failure of vancomycin oral.
Age Restrictions	6 months or older
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Drizalma (duloxetine)

Products Affected

- Drizalma Sprinkle

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting a trial or failure of duloxetine (Cymbalta) capsule or amitriptyline use in the last 180 days.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dupixent

Products Affected

- Dupixent

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Atopic Dermatitis a. Diagnosis of moderate to severe atopic dermatitis. b. 6 months of age and older c. Trial and failure, contraindication, or intolerance to a medium to high potency topical steroid (e.g. mometasone, fluocinolone, fluocinonide). d. IGA score of at least 3 e. EASI score of at least 16 f. Minimum body surface area involvement of =10% g. Initial curation: 6 months h. Reauthorization: Documentation of positive clinical response and will not be used in combination with another biologic medication.</p> <p>2. Asthma a. Diagnosis of moderate to severe asthma defined as pre-bronchodilator FEV1 =80% b. 6 years of age and older c. History of one or more asthma exacerbations that required treatment with systemic corticosteroids or emergency visit or hospitalization for the treatment of asthma within the past year. d. Daily dependence on oral corticosteroids in addition to regular use of high-dose inhaled corticosteroids plus an additional controller. e. Blood eosinophils =300 cells/mcL f. Initial coverage duration: 6 months g. Reauthorization: Documented clinical response to Dupixent demonstrated by 1) reduction in frequency of exacerbations, 2) decreased utilization of rescue medications, 3) increase in FEV1 from pretreatment baseline, 4) reduction in oral corticosteroid requirements: Dupixent will be used in combination with an ICS controller medication: and Dupixent will not be used with another biologic medication.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Continuation: 6 months

PA Criteria	Criteria Details
Other Criteria	<p>3. Chronic Rhinosinusitis with Nasal Polyps</p> <p>a. 18 years of age and older</p> <p>b. To be used as add-on maintenance treatment for individuals with:</p> <ul style="list-style-type: none"> • Nasal polyps detected by direct examination, endoscopy, or sinus CT scan • Significant rhinosinusitis such as nasal obstruction, rhinorrhea, or reduction or loss of smell as documented by the prescriber. <p>c. Bilateral Nasal Polyp Score (NPS) of at least 5, and NPS of at least 2 in each nostril.</p> <p>d. Documented inadequate response to nasal corticosteroids.</p> <p>e. Patient has received treatment with systemic corticosteroids with the past two years (or has a contraindication) or has had prior surgery for nasal polyps.</p> <p>f. Initial coverage: 6 months</p> <p>g. Reauthorization: Documented positive clinical response: patient will continue to receive Dupixent as add-on maintenance therapy in combination with intranasal corticosteroids, and patient will not use Dupixent with another biologic medication.</p> <p>4. Eosinophilic Esophagitis:</p> <p>a. Documented trial and failure of a proton pump inhibitor (PPI) or a topical glucocorticoid steroid.</p> <p>b. Diagnosis confirmed by greater than or equal to 15 intraepithelial eosinophils per high-power field (eos/hpf).</p> <p>c. Prescribed by or in consultation with a gastroenterologist or allergist.</p> <p>d. Reauthorization: documented positive clinical response as demonstrated by a decrease in eos/hpf and improvement in baseline Dysphagia Symptom Questionnaire (DSQ) score.</p> <p>5. Prurigo Nodularis:</p> <p>a. Worst Itch-Numeric Rating Scale (WI-NRS) greater than or equal to 7 and 20 or more nodular lesions.</p> <p>b. Inadequate response, intolerance, or contraindication to a high potency topical steroid (e.g., betamethasone, fluocinonide, triamcinolone).</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Emend (aprepitant oral)

Products Affected

- Aprepitant Oral Capsule
- Emend Oral Suspension Reconstituted

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Emgality (galcanezumab-gnlm)

Products Affected

- Emgality

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Coverage for a Diagnosis of Episodic Migraines and Chronic Migraines requires a History of failure or intolerance to two (2) preventive migraine medications from at least two (2) of the following classes: anti-depressant (e.g. venlafaxine), anti-convulsant (e.g. topiramate, divalproex) or anti-hypertensive (e.g. propranolol, verapamil). Coverage for a diagnosis of Episodic Cluster headaches requires a History of failure or intolerance to Verapamil.
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with a Headache Specialist, a Neurologist, or a Pain Specialist.
Coverage Duration	6 months
Other Criteria	Continuation: Documentation that the patient has experienced a positive clinical response
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Enbrel (etanercept)

Products Affected

- Enbrel Mini
- Enbrel Subcutaneous Solution 25 MG/0.5ML
- Enbrel Subcutaneous Solution Prefilled Syringe
- Enbrel Subcutaneous Solution Reconstituted
- Enbrel SureClick Subcutaneous Solution Auto-Injector

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1) Ankylosing Spondylitis (AS) - Trial and failure of NSAID. Patients with peripheral arthritis must have a trial of sulfasalazine. Patients with axial disease and failure of NSAIDs can be started on Enbrel without a trial of sulfasalazine. 2) Juvenile Idiopathic Arthritis (JIA) An adequate trial (3 months or more) of one of the following DMARDs: i. Leflunomide ii. Methotrexate iii. Sulfasalazine. 3) Psoriatic Arthritis (PsA) at least a three months trial of one of the following: cyclosporine, leflunomide (LEF), MTX, SSZ 4) Rheumatoid Arthritis (RA) - Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. c. An adequate trial (3 months or more) of one of the following other DMARDs: i. Hydroxychloroquine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine 5) Plaque Psoriasis (PsO) 1. Plaque Psoriasis (PsO): a. The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to, or is intolerant to a 3-month trial of a topical agent (topical corticosteroid, topical calcineurin inhibitor, topical vitamin D analogs, etc.).</p>
Age Restrictions	
Prescriber Restrictions	<p>1. AS- prescribed by or in consultation with a rheumatologist 2. JIA- prescribed by or in consultation with a rheumatologist 3. PsA- prescribed by or in consultation with a dermatologist or rheumatologist 4. PsO-</p>

PA Criteria	Criteria Details
	prescribed by or in consultation with a dermatologist 5. RA- prescribed by or in consultation with a rheumatologist
Coverage Duration	1 year
Other Criteria	For all diagnoses: 1. Current PPD (tuberculosis) negative skin test or negative QuantiFERON-TB Gold test before the initiation of therapy, 2. The appropriate Disease Specific Criteria has been met. CONTINUATION CRITERIA: Documentation of positive response with Enbrel treatment
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Endari (L-glutamine)

Products Affected

- Endari

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis of sickle cell disease. 2. Will be used to reduce acute complications of sickle cell disease. 3. Will be used concurrently with hydroxyurea, unless contraindicated or not tolerated. 4. Member has had two (2) or more painful sickle cell crises within the past 12 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	Continuation therapy: positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Entocort (budesonide capsules)

Products Affected

- Budesonide Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. The member has a diagnosis of mild to moderate Crohn's disease. 2. Documented trial and failure of sulfasalazine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epclusa (sofosbuvir/velpatasvir)

Products Affected

- Sofosbuvir-Velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical records documenting the diagnosis of chronic Hepatitis C, including laboratory documentation of genotype and subtype, detectable HCV RNA levels at baseline, HIV status and liver transplant status.
Age Restrictions	3 years of age or older
Prescriber Restrictions	Gastroenterologist or Infectious Disease specialist or Hepatologist
Coverage Duration	Duration as per package insert or Class I or II recommendation by the AASLD/IDSA/IAS-USA guidelines
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epidiolex (cannabidiol)

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Lennox-Gastaut Syndrome: Diagnosis of Lennox-Gastaut Syndrome and seizures have been inadequately controlled by a trial of at least two antiepileptic drugs (e.g. clobazam, valproate, lamotrigine, topiramate, levetiracetam). Dravet Syndrome: Diagnosis of Dravet Syndrome and seizures have been inadequately controlled by a trial of at least two antiepileptic drugs (e.g. clobazam, valproate, lamotrigine, topiramate, levetiracetam).
Age Restrictions	1 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a Neurologist
Coverage Duration	Initial: 6 months ReAuthorization: 1 year
Other Criteria	AST, ALT and total bilirubin levels will be obtained 1 month, 3 months, and 6 months after initiation, then periodically thereafter or as clinically indicated. ReAuthorization: Patient is tolerating treatment and there continues to be a medical need for the medication and there has been disease stabilization or improvement while on this medication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Erivedge (vismodegib)

Products Affected

- Erivedge

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Erleada (apalutamide)

Products Affected

- Erleada

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Erythropoietin Stimulating Agents

Products Affected

- Retacrit Injection Solution 10000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 40000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Esbriet (pirfenidone)

Products Affected

- Pirfenidone Oral Capsule
- Pirfenidone Oral Tablet 267 MG, 801 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Patient must have a Forced Vital Capacity (FVC) greater than or equal to 50% of predicted.
Age Restrictions	18 years or older
Prescriber Restrictions	Pulmonologist
Coverage Duration	1 year
Other Criteria	Baseline Liver Function test
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Exkivity

Products Affected

- Exkivity

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Extavia (interferon beta-1b)

Products Affected

- Extavia Subcutaneous Kit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Trial and failure of Avonex (interferon beta 1a) and Rebif (interferon beta 1a)
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Farydak (panobinostat)

Products Affected

- Farydak

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or there must be a Class I or IIa recommendation in the Thomson Micromedex DrugDex compendium.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fasenra (benralizumab)

Products Affected

- Fasenra Pen
- Fasenra Subcutaneous Solution Prefilled Syringe 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FDA approved for asthma maintenance add-on therapy in severe asthma (eosinophilic phenotype), medical records must document IgE level greater than or equal to 150 cells/microliter, AND inadequate control with an inhaled corticosteroid and a long acting beta-2 agonist combination therapy, AND evidence of persistent symptoms requiring frequent rescue therapy, practitioner visits despite inhaled corticosteroids, ER visits OR inadequate control OR intolerance OR contraindication to inhaled corticosteroid and a long acting beta-2 agonist combination. Not to be used as monotherapy or concomitantly with other biologics. Continuation Criteria: Documentation of a reduction in exacerbation frequency and/or severity.
Age Restrictions	
Prescriber Restrictions	Allergist, Pulmonologist, Dermatologist or Immunologist
Coverage Duration	Initial: 9 months Continuation: 12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fensolvi (leuprolide)

Products Affected

- Fensolvi
- Fensolvi (6 Month)

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for treatment of prostate cancer
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ferriprox (deferiprone)

Products Affected

- Deferiprone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Patient must have an absolute neutrophil count (ANC) greater than $1.5 \times 10^9/L$ before starting therapy. Patient must have tried and failed or have a contraindication to deferasirox. ANC levels are not required for iron transfusional overload in patients with sickle cell disease or other anemias.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For renewal, must receive documentation demonstrating clinical efficacy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Firazyr (icatibant)

Products Affected

- Icatibant Acetate
- Sajazir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical records must document a diagnosis of hereditary angioedema (HAE) based on evidence of a low C4 level and one of the following: a low C1 inhibitor (C1-INH) antigenic level or a normal C1-INH antigenic level and a low C1-INH functional level. The patient must not be concurrently taking an angiotensin converting enzyme (ACE) inhibitor or estrogen replacement therapy and must be experiencing at least one symptom of a moderate or severe attack (ie. swelling of the face, throat, or abdomen).
Age Restrictions	18 years or older
Prescriber Restrictions	Allergist or Immunologist
Coverage Duration	6 months
Other Criteria	Medical records documenting frequency of acute HAE attacks and the patient's response to therapy must be provided. If the patient is experiencing more than one acute HAE attack per month medical records documenting use of a long-term prophylactic therapy (LTP) or the clinical rationale for avoiding LTP must be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Firdapse (amifampridine)

Products Affected

- Firdapse

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Confirmed diagnosis of Lambert-Eaton myasthenic syndrome (LEMS), documentation of baseline confirmatory diagnostic test results including but not limited to: a Repetitive Nerve Stimulation (RNS), a positive anti-P/Q type voltage-gated calcium channel antibody test, a Quantitative Myasthenia Gravis (QMG) score, a triple-timed up-and-go test (3TUG), a Timed 25-foot walk test (T25FW).
Age Restrictions	6 years of age and older
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 3 months Renewal: 6 months
Other Criteria	Dose does not exceed 80mg per day. Renewal: Documentation of clinical improvement in symptoms
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Flexeril (cyclobenzaprine)

Products Affected

- Cyclobenzaprine HCl Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Prior authorization required for patients 65 years of age or older.
Age Restrictions	Prior authorization required for patients 65 years of age or older.
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Forteo (teriparatide)

Products Affected

- Forteo Subcutaneous Solution 600 MCG/2.4ML
- Forteo Subcutaneous Solution Pen-Injector
- Teriparatide (Recombinant) Subcutaneous Solution Pen-Injector 620 MCG/2.48ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	A diagnosis of osteoporosis and a T-score of -2.5 or less at the femoral neck, total hip or lumbar spine by DXA. The patient has failed or is intolerant to other available osteoporosis therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fotivda (Tivozanib)

Products Affected

- Fotivda

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Fruzaqla (fruquintinib)

Products Affected

- Fruzaqla

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Gamunex-C (human immune globulin)

Products Affected

- Flebogamma DIF
- Gamunex-C

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gattex (teduglutide)

Products Affected

- Gattex

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Patient must be dependent on parenteral nutrition/intravenous (PN/I.V.) support for at least 12 months and requires PN at least 3 times per week.
Age Restrictions	1 year of age and older
Prescriber Restrictions	Gastroenterologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gavreto (pralsetinib)

Products Affected

- Gavreto

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Gilotrif (afatinib)

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Gleevec (imatinib)

Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GLP-1 Agonists

Products Affected

- Mounjaro
- Ozempic (0.25 or 0.5 MG/DOSE)
- Ozempic (1 MG/DOSE) Subcutaneous Solution Pen-Injector 4 MG/3ML
- Ozempic (2 MG/DOSE)
- Rybelsus
- Trulicity
- Victoza Subcutaneous Solution Pen-Injector

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis of type 2 diabetes. 2. A 90-day prescription fill history of metformin within the past 180 days.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hadlima (adalimumab-bwwd)

Products Affected

- Hadlima
- Hadlima PushTouch

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Ankylosing Spondylitis (AS): peripheral arthritis must have a trial of sulfasalazine and an NSAID. Patients with axial disease and failure of NSAIDs can be started without a trial of sulfasalazine. 2. Crohn's disease (CD): Inadequate response or intolerance to conventional therapy. For the purpose of this policy, conventional therapy includes the use of ONE of the following: i. Corticosteroids (e.g., prednisone, prednisolone, dexamethasone, budesonide). ii. Methotrexate iii. Thiopurines (azathioprine, mercaptopurine). 3. Juvenile Idiopathic Arthritis (JIA): An adequate trial (3 months or more) of one of the following DMARDs: i. Leflunomide ii. Methotrexate iii. Sulfasalazine. 4. Plaque Psoriasis (PsO): a. The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to, or is intolerant, a 3-month trial of a topical agent (topical corticosteroid, topical calcineurin inhibitor, topical vitamin D analogs, etc.). 5. Psoriatic Arthritis (PsA): at least a three month trial of one of the following: cyclosporine, leflunomide (LEF), MTX, SSZ 6. Rheumatoid Arthritis (RA): Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. c. An adequate trial (3 months or more) of one of the following other DMARDs: i. Hydroxychloroquine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine 7. Ulcerative Colitis (UC): The member has had an inadequate response to one of the following: aminosalicylates, corticosteroids, thiopurines, or cyclosporine.</p>
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist, dermatologist, gastroenterologist, or ophthalmologist.

PA Criteria	Criteria Details
Coverage Duration	One (1) year
Other Criteria	8. Hidradenitis Suppurativa (HS): Documented diagnosis of Hurley Stage III HS or refractory Hurley Stage II hidradenitis suppurativa. 9.Uveitis: a. Documented diagnosis of non-infectious intermediate, posterior and panuveitis in adult patients and meets the following: i. A documented trial and failure, contraindication, or intolerance to conventional therapy such as ophthalmic or systemic corticosteroids AND immunosuppressive drugs (e.g. azathioprine, cyclosporine, methotrexate, or tacrolimus). For all diagnoses: 1. Current PPD (tuberculosis) negative skin test or negative QuantiFERON-TB Gold test before the initiation of therapy, 2. The appropriate Disease Specific Criteria has been met. CONTINUATION CRITERIA: Documentation of positive response with treatment
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hetlioz (tasimelteon)

Products Affected

- Tasimelteon

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Must have a documented diagnosis of (1) Non-24-Hour Sleep-Wake Disorder in adults or (2) Nighttime sleep disturbances in Smith-Magenis Syndrome (SMS) in patients 16 years of age and older.
Age Restrictions	
Prescriber Restrictions	Sleep Specialist or Neurologist
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	For renewal, chart notes must show clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hexalen (altretamine)

Products Affected

- Hexalen

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or there must be a Class I or IIa recommendation in the Thomson Micromedex DrugDex compendium.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High Risk Medication - Antidepressants TCA

Products Affected

- Amitriptyline HCl Oral
- Amoxapine
- clomiPRAMINE HCl Oral
- Desipramine HCl Oral
- Doxepin HCl Oral Capsule
- Imipramine HCl Oral
- Nortriptyline HCl Oral Capsule
- Nortriptyline HCl Oral Solution
- Protriptyline HCl
- Trimipramine Maleate Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	No Prior Authorization is required if 64 years of age or less.
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High Risk Medication - Paroxetine

Products Affected

- PARoxetine HCl
- Paxil Oral Suspension
- PARoxetine HCl ER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	You must have taken two (2) of the following drugs: a selective serotonin reuptake inhibitor (SSRI) or a serotonin norepinephrine reuptake inhibitor (SNRI).
Age Restrictions	No Prior Authorization is required if 64 years of age or less.
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High Risk Medications

Products Affected

- chlordiazePOXIDE HCl
- chlorproMAZINE HCl Oral Tablet
- Scopolamine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	No Prior Authorization is required if 64 years of age or less.
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High Risk Medications (estrogen containing products)

Products Affected

- Alora
- Amabelz
- CombiPatch
- Dotti
- Estradiol Oral Tablet 2 MG
- Estradiol-Norethindrone Acet
- Estropipate Oral
- Evamist
- Fyavolv
- Jevantique Lo
- Jinteli
- Lopreeza
- Lyllana
- Menest
- Mimvey
- Mimvey Lo
- Norethindrone-Eth Estradiol
- Prefest
- Premarin Oral
- Premphase
- Prempro

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	No Prior Authorization is required if 64 years of age or less.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Ibrance (palbociclib)

Products Affected

- Ibrance

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Iclusig (ponatinib)

Products Affected

- Iclusig

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Idhifa (enasidenib)

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	Documentation of an isocitrate dehydrogenase-2 (IDH2) mutation detected by an FDA-approved test.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Imbruvica (ibrutinib)

Products Affected

- Imbruvica Oral Capsule
- Imbruvica Oral Suspension
- Imbruvica Oral Tablet 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Increlex (mecasermin)

Products Affected

- Increlex

PA Criteria	Criteria Details
Exclusion Criteria	Secondary forms of IGF-1 deficiency.
Required Medical Information	Documentation of all of the following: height standard deviation score less than or equal to negative 3, basal IGF-1 standard deviation score less than or equal to negative 3, normal or elevated GH levels, predicted adult height more than 1.5 standard deviations below the mid-parenteral height, growth rate less than 7 cm/year if less than or equal to 3 years old and less than 5 cm if greater than 3 years old, and open epiphyses on bone radiograph.
Age Restrictions	2 to 18 years of age
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inlyta (axitinib)

Products Affected

- Inlyta

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Inqovi (cedazuridine/decitabine)

Products Affected

- Inqovi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Inrebic (fedratinib)

Products Affected

- Inrebic

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Iressa (gefitinib)

Products Affected

- Gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Ivermectin (s)

Products Affected

- Ivermectin Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Strongyloidiasis: Diagnosis of intestinal (i.e., nondisseminated) strongyloidiasis due to the nematode parasite Strongyloides stercoralis OR both of the following: member received the drug within the past 120 days and member requires continuation of therapy. Onchocerciasis: Diagnosis of onchocerciasis due to the nematode parasite Onchocerca volvulus OR both of the following: member received the drug within the past 120 days and member requires continuation of therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Strongyloidiasis: 3 weeks. Onchocerciasis: 6 months.
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Iwilfin (eflornithine)

Products Affected

- Iwilfin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Jakafi (ruxolitinib)

Products Affected

- Jakafi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Jaypirca (pirtobrutinib)

Products Affected

- Jaypirca

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomsn Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Juxtapid (lomitapide)

Products Affected

- Juxtapid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	A diagnosis of homozygous familial hypercholesterolemia (HoFH) as defined by the presence of at least one of the following clinical criteria: (1) documented functional mutation(s) in both low-density lipoprotein (LDL) receptor alleles or alleles known to affect LDL receptor functionality, or (2) skin fibroblast LDL receptor activity less than 20% normal, or (3) untreated total cholesterol (TC) greater than 500 mg/dL and triglycerides (TG) less than 300 mg/dL and both parents with documented untreated TC greater than 250 mg/dL. Patient must have tried and failed a high intensity statin at maximum tolerated dose OR have a documented intolerance or contraindication to statins. Must be used in combination with other lipid-lowering treatments.
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jylamvo (methotrexate)

Products Affected

- Jylamvo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Kalydeco (ivacaftor)

Products Affected

- Kalydeco Oral Packet 25 MG, 50 MG, 75 MG
- Kalydeco Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	1 month old to 5 years of age: Oral granules. 6 years of age or older: Oral tablets.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kerendia (finerenone)

Products Affected

- Kerendia

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Member has a documented diagnosis chronic kidney disease (CKD) associated with type 2 diabetes (T2D). 2. Diagnosis is defined by one of the following: a. UACR of 30 to 300 mg/g, minimum eGFR of 25 mL/min/1.73 m², and diabetic retinopathy, or, b. UACR of greater than or equal to 300 mg/g and a minimum eGFR 25 ml/min/1.73 m². 3. Therapy will not be initiated if serum potassium is greater than 5 mEq/L. 4. Member has been on a maximally tolerated dose of either an angiotensin converting enzymy (ACE) inhibitor or angiotensin receptor blocker (ARB), unless otherwise contraindicated. 5. Mmeber has tried and had an inadequate response or intolerance to a sodium-glucose cotransporter (SGLT2) inhibitor (e.g., Farxiga).</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	Continuation criteria: Documentation of positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Kineret (anakinra)

Products Affected

- Kineret Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Exclusion Criteria	Will not be approved for use in combination with TNF antagonists.
Required Medical Information	Rheumatoid Arthritis (RA): Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. An adequate trial (3 months or more) of one of the following other DMARDs: hydroxychloroquine, leflunomide, methotrexate, sulfasalazine. Trial and failure of two of the following, unless contraindicated or not tolerated: Amjevita, Enbrel, Hadlima, Rinvoq, Xeljanz.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	Continuation criteria: positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kisqali (ribociclib)

Products Affected

- Kisqali (200 MG Dose)
- Kisqali (400 MG Dose)
- Kisqali (600 MG Dose)
- Kisqali 200 Dose
- Kisqali 400 Dose
- Kisqali 600 Dose
- Kisqali Femara (200 MG Dose)
- Kisqali Femara (400 MG Dose)
- Kisqali Femara (600 MG Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Korlym (mifepristone)

Products Affected

- miFEPRIStone Oral Tablet 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Koselugo (selumetinib)

Products Affected

- Koselugo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Krazati (adagrasib)

Products Affected

- Krazati

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Kuvan (sapropterin)

Products Affected

- Sapropterin Dihydrochloride

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Initial Authorizations: Medical records documenting all of the following: the target Phe blood level for the patient and dose does not exceed FDA approved maximum for the diagnosis. Renewal: Recent Phe level is at target range.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial authorization: 2 months Renewal: 6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lenvima (lenvatinib)

Products Affected

- Lenvima (10 MG Daily Dose)
- Lenvima (12 MG Daily Dose)
- Lenvima (14 MG Daily Dose)
- Lenvima (18 MG Daily Dose)
- Lenvima (20 MG Daily Dose)
- Lenvima (24 MG Daily Dose)
- Lenvima (4 MG Daily Dose)
- Lenvima (8 MG Daily Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Leukine (sargramostim)

Products Affected

- Leukine Injection Solution Reconstituted
- Leukine Intravenous

PA Criteria	Criteria Details
Exclusion Criteria	Combination use with pegfilgrastim or filgrastim.
Required Medical Information	Medical records documenting neutropenia which is defined as an absolute neutrophil count less than 500/mm ³ .
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lidoderm (lidocaine topical patch)

Products Affected

- Lidocaine External Patch 5 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lonsurf (tipiracil/trifluridine)

Products Affected

- Lonsurf

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lorbrena (lorlatinib)

Products Affected

- Lorbrena

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lotronex (alosetron)

Products Affected

- Alosetron HCl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis of severe diarrhea-predominant irritable bowel syndrome (IBS) with symptoms persisting for at least six (6) months. 2. Patient was female at birth. 3. Patient has not responded adequately to conventional therapy (i.e., loperamide, antispasmodics).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lumakras (sotorasib)

Products Affected

- Lumakras

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lybalvi

Products Affected

- Lybalvi

PA Criteria	Criteria Details
Exclusion Criteria	Know opioid use disorder or is dependent on opioids for a chronic health condition. Patients undergoing acute opioid withdrawal.
Required Medical Information	Patient is 18 years of age or older. Patient has a diagnosis of schizophrenia or bipolar I disorder made by a behavior health practitioner or in consultation with a behavior health practitioner. Patient does not have a known opioid use disorder or is dependent on opioids for a chronic health condition. There should be a 7-day opioid-free interval from the last use of short-acting opioids, and at least a 14-day opioid-free interval from the last use of long-acting opioids to avoid precipitation of opioid withdrawal. Trial of generic olanzapine with documentation demonstrating positive therapeutic benefit but unacceptable weight gain while on therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lynparza (olaparib)

Products Affected

- Lynparza

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Lyrica CR (pregabalin)

Products Affected

- Pregabalin ER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Trial and failure of gabapentin and Lyrica capsule
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lytgobi (futibatinib)

Products Affected

- Lytgobi (12 MG Daily Dose)
- Lytgobi (16 MG Daily Dose)
- Lytgobi (20 MG Daily Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomsn Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Marinol (dronabinol)

Products Affected

- Dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	If using for nausea and vomiting associated with cancer chemotherapy, the patient must be receiving cancer chemotherapy and has failed one 5-HT3 antagonist and one of the following: corticosteroid, anti-histamine, anti-psychotic, or prokinetic. If using for anorexia associated with weight loss due to HIV/AIDS, the patient must have a trial and failure, contraindication or intolerance to Megace ES (megestrol oral suspension).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Matulane (procarbazine)

Products Affected

- Matulane

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Mavyret (glecaprevir/pibrentasvir)

Products Affected

- Mavyret

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical records documenting the diagnosis of chronic Hepatitis C, including laboratory documentation of genotype and subtype, detectable HCV RNA levels at baseline, HIV status and liver transplant status.
Age Restrictions	3 year of age and older.
Prescriber Restrictions	Gastroenterologist or Infectious Disease specialist or Hepatologist
Coverage Duration	Duration as per package insert or Class I or II recommendation by the AASLD/IDSA/IAS-USA guidelines
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mekinist (trametinib)

Products Affected

- Mekinist

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MEKTOVI (binimetinib)

Products Affected

- Mektovi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Movantik (naloxegol)

Products Affected

- Movantik

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting a trial/failure of lactulose oral or lubiprostone.
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Natpara (parathyroid hormone)

Products Affected

- Natpara

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical records must document a diagnosis of hypocalcemia due to chronic hypoparathyroidism. The prescriber must provide documentation that the patient must have a normal 25-hydroxyvitamin D level and a serum calcium level above 7.5mg/dL. For renewal the albumin-corrected total serum calcium level between 7.5 mg/dL and 10.6 mg/dL should be achieved with standard of care.
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nerlynx (neratinib)

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	Lab values for total bilirubin, AST, ALT, and alkaline phosphate levels prior to starting treatment.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Nexavar (sorafenib)

Products Affected

- SORafenib Tosylate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Ninlaro (ixazomib)

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Northera (droxidopa)

Products Affected

- Droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	2 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Noxafil (posaconazole)

Products Affected

- Posaconazole Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	2 years of age and older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nubeqa (darolutamide)

Products Affected

- Nubeqa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Nuedexta (dextroamphetamine / quinidine)

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting a neurological disease or injury. Chart notes documenting a diagnosis of Pseudobulbar Affect (PBA) secondary to a neurological disease or injury (e.g. Multiple Sclerosis, ALS, Parkinson's, stroke, traumatic brain injury). Documentation of the number of PBA episodes per day. Chart notes documenting medications tried/failed to reduce the number of PBA episodes. If the prescriber is not a specialist then the referral notes of the specialist must be submitted. Baseline Center for Neurologic Study-Lability Scale (CNS-LS) score of 13 or greater.
Age Restrictions	
Prescriber Restrictions	Must be prescribed by or in consultation with a neurologist.
Coverage Duration	Initial: 3 months Continuation of therapy: 6 months
Other Criteria	Continuation of Treatment Criteria: medical records documenting the following: 1. A decrease in the CNS-LS score and the decrease has been maintained. 2. A decrease in the number of daily episodes.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuplazid (pimavanserin)

Products Affected

- Nuplazid Oral Capsule
- Nuplazid Oral Tablet 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	For renewal: Chart notes must document an improvement in psychosis symptoms from baseline and demonstrates a continued need for treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuvigil (armodafinil)

Products Affected

- Armodafinil

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	If using for obstructive sleep apnea (OSA), must provide documentation that the patient has OSA and has been evaluated by a sleep specialist.
Age Restrictions	
Prescriber Restrictions	Neurologist or Sleep Specialist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ocrevus (ocrelizumab)

Products Affected

- Ocrevus

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting a diagnosis of Primary Progressive Multiple Sclerosis. Patients with primary progressive multiple sclerosis will not be required to have a trial and failure of other medications. Diagnosis of Relapsing Multiple Sclerosis, chart notes must document a trial/failure of interferon or glatiramer.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Odomzo (sonidegib)

Products Affected

- Odomzo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Ofev (nintedanib)

Products Affected

- Ofev

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	18 years or older
Prescriber Restrictions	Pulmonologist
Coverage Duration	1 year
Other Criteria	Baseline Liver Function tests
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ogsiveo (nirogacestat)

Products Affected

- Ogsiveo Oral Tablet 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6 months)
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Ojjaara (momelotinib)

Products Affected

- Ojjaara

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Omnitrope (somatropin)

Products Affected

- Omnitrope

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Adult onset growth hormone deficiency (GHD) - Multiple hormone deficiencies (hypopituitarism) resulting from pituitary disease, hypothalamic disease, surgery, radiation therapy or trauma. a) Patient has greater than or equal to 2 of the following pituitary hormone deficiencies: thyroid stimulating hormone deficiency, adrenocorticotropin hormone deficiency, gonadotropin deficiency, an arginine vasopressin (AKA vasopressin or antidiuretic hormone (ADH)) deficiency. b) Low serum IGF-I. c) Patient must exhibit clinical features of adult GHD including: increased body fat, decreased muscle mass, poor exercise performance, decreased bone density, and cardiovascular risk factors (high LDL, low HDL). d) Documentation of baseline information (IGF-I levels, lipids, bone density, cardiovascular factors, body composition, exercise capacity) provided with each request. Childhood onset GHD - Adults who were GH deficient as children or adolescents. a) Patient has subnormal response to at least 2 provocative stimulation tests (less than or equal to 5 ng/ml) following a GH washout period of 1-3 months. b) Patient must exhibit clinical features of adult GHD including: increased body fat, decreased muscle mass, poor exercise performance, decreased bone density, and cardiovascular risk factors. c) Documentation of baseline information (IGF-I levels, lipids, bone density, cardiovascular factors, body composition, exercise capacity) provided with each request.</p>
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	1 year
Other Criteria	

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Onureg (Azacitidine)

Products Affected

- Onureg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Opsumit (macitentan)

Products Affected

- Opsumit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Patient must be classified as WHO functional class (FC) II-IV and must have tried and failed or have a contraindication to Revatio (sildenafil) or Adcirca (tadalafil).
Age Restrictions	18 years or older
Prescriber Restrictions	Cardiologist or Pulmonologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orencia (abatacept SubQ)

Products Affected

- Orencia ClickJect
- Orencia Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with TNF antagonists.
Required Medical Information	Rheumatoid Arthritis (RA): Moderate to severe RA is defined as: DAS-28 more than 3.2 or CDAI more than 10.1. An adequate trial (3 months or more) of one of the following other DMARDs: hydroxychloroquine, leflunomide, methotrexate, sulfasalazine. Inadequate response, unless contraindicated or not tolerated to two of the following: Amjevita, Enbrel, Hadlima, Rinvoq, Xeljanz.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Continuation: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orfadin (nitisinone)

Products Affected

- Nitisinone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orgovyx (Relugolix)

Products Affected

- Orgovyx

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Orkambi (ivacaftor/lumacaftor)

Products Affected

- Orkambi Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	1 year and older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orserdu (elacestrant)

Products Affected

- Orserdu

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomsn Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Otezla (apremilast)

Products Affected

- Otezla

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Psoriatic Arthritis, active (PsA): The patient has had at least a trial and failure of two of the following: Amjevita, Cosentyx, Enbrel, Hadlima, Rinvoq, Skyziri, or Xeljanz. Plaque psoriasis, moderate to severe (PsO): The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to two of the following: Amjevita, Cosentyx, Enbrel, Hadlima, or Skyrizi. Behcet's syndrome: trial and failure of adalimumab (Amjevita)
Age Restrictions	Approved for patients 18 years of age or older
Prescriber Restrictions	Must be prescribed by or in consultation with a Rheumatologist, Dermatologist or Ophthalmologist
Coverage Duration	Initial: 6 months Reauthorization: 1 year
Other Criteria	Reauthorization: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pemazyre (pemigatinib)

Products Affected

- Pemazyre

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Perseris (risperidone)

Products Affected

- Perseris

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a trial/failure of Risperdal Consta.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Piqray (alpelisib)

Products Affected

- Piqray (200 MG Daily Dose)
- Piqray (250 MG Daily Dose)
- Piqray (300 MG Daily Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Pomalyst (pomalidomide)

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Prolastin (alpha-1-proteinase inhibitor [human])

Products Affected

- Prolastin-C

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of congenital alpha1-antitrypsin deficiency with clinically evident emphysema and chronic replacement therapy is needed. The member has an alpha1-antitrypsin phenotype of PiZZ, PiZ(null), or Pi (null, null) or phenotypes associated with serum alpha 1-antitrypsin concentrations of less than 50mg/dL if/when measured by laboratories using nephelometry instead of radial immunodiffusion. Otherwise, a deficiency is shown at 80mg/dL. (These products should not be used in individuals with the PiMZ or PiMS phenotypes of alpha1-antitrypsin deficiency because these individuals appear to be at small risk of developing clinically evident emphysema.)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Continuation: 1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Promacta (eltrombopag olamine)

Products Affected

- Promacta

PA Criteria	Criteria Details
Exclusion Criteria	Coverage will not be provided when used in combination with Nplate.
Required Medical Information	If using for idiopathic thrombocytopenia, patient must have a trial and failure or contraindication to at least two of the following treatments: corticosteroids, immunoglobulin, or splenectomy. Dose may not exceed the FDA-approved maximum dose. If using for thrombocytopenia due to chronic hepatitis C, and the patient is currently on antiviral interferon therapy or will be starting interferon therapy, the patient must have a platelet count less than 75,000/microliter. Severe aplastic anemia (initial): Diagnosis of severe aplastic anemia. Patient has a platelet count less than 30,000/microliter. Trial and failure, intolerance, or contraindication to immunosuppressive therapy with antithymocyte globulin and cyclosporine. First-line treatment of severe aplastic anemia in combination with standard immunosuppressive therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pulmonary Arterial Hypertension - ERA

Products Affected

- Ambrisentan
- Bosentan

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, Idiopathic Pulmonary Fibrosis
Required Medical Information	1. Member has a confirmed diagnosis of primary or secondary pulmonary arterial hypertension (WHO Group 1) by right heart catheterization. 2. Confirmed diagnosis will show all of the following: a) Mean pulmonary artery pressure (mPAP) greater than 20 mmHG at rest, b) Pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg, c) Pulmonary vascular resistance greater than 3 Wood units. 3. Individual has WHO functional class II-IV symptoms.
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	Initial: 6 months Continuation: 1 year
Other Criteria	Continuation criteria: Chart notes documentating clinically significant improvement or stabilization in clinical signs and symptoms of disease (including but not limited to improvement in walk distance, dyspnea, and/or functional class).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Purixan (mercaptopurine)

Products Affected

- Purixan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Qinlock (ripretinib)

Products Affected

- Qinlock

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Qualaquin (quinine sulfate)

Products Affected

- QuiNINE Sulfate Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Regranex (becaplermin)

Products Affected

- Regranex

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation confirming treatment is or lower extremity diabetic neuropathic ulcers that extend in to the subcutaneous tissue and have adequate supply. This medication will be used as an adjunct to, and not a substitute for, good ulcer care practices including initial sharp debridement, pressure relief, and infection control.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Relistor (methylnaltrexone)

Products Affected

- Relistor Oral
- Relistor Subcutaneous Solution

PA Criteria	Criteria Details
Exclusion Criteria	Non-opioid induced constipation. Fecal impaction. Acute diverticular disease. Acute surgical abdomen.
Required Medical Information	Documentation that the patient is on chronic opioid therapy, and documented trial and failure of naloxegol (for approval of oral tablets only). Documented trial and failure of naloxegol not required for injection.
Age Restrictions	18 years or older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Repatha (evolocumab)

Products Affected

- Repatha
- Repatha SureClick
- Repatha Pushtronex System

PA Criteria	Criteria Details
Exclusion Criteria	Homozygous Familial Hypercholesterolemia (HoFH) for members with two LDL receptor negative alleles. Will not be approved for use in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor.
Required Medical Information	Chart notes documenting one of the following diagnoses: 1) Atherosclerotic cardiovascular disease (ASCVD) (e.g. Acute coronary syndrome, history of myocardial infarction, angina, arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin) OR 2) Heterozygous Familial Hypercholesterolemia (HeFH) OR 3) Homozygous Familial Hypercholesterolemia (HoFH), OR 4) Primary Hyperlipidemia. For HeFH: results of Dutch Lipid Clinic diagnostic criteria score greater than or equal to 9. For HoFH: results of a genetic test OR an untreated LDL level over 500mg/dl AND the presence of xanthoma before 10 years old or evidence of HeFH in both parents AND 1 of the following: 1) Member has received at least 3 months of high intensity (HI) statin at maximum tolerated dose (MTD), OR 2) Member can't tolerate HI statin and has received 3 months of a moderate intensity statin or a low intensity statin at a MTD, OR 3) Member is unable to tolerate two statin medications, one of which is HI OR 4) Member has a contraindication to all statins documented in chart notes, OR 5) Member has experienced rhabdomyolysis. Member has been on at least 3 months of ezetimibe therapy as adjunct to MTD of statin therapy and will continue to receive it OR Member has history of, failure, contraindication or intolerance to ezetimibe. Chart notes within the past 30 days documenting baseline and a current lipid panel while on maximum tolerated lipid lowering regimen.
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with one of the following: cardiologist, endocrinologist, or lipid specialist.

PA Criteria	Criteria Details
Coverage Duration	Initial: 3 months, Continuation: 6 months
Other Criteria	High intensity statins: atorvastatin 40-80mg, rosuvastatin 20-40mg. Moderate intensity statins: atorvastatin 10-20mg, rosuvastatin 5-10mg, simvastatin 20mg or more, pravastatin 40mg or more, lovastatin 40mg, fluvastatin XL 80mg, fluvastatin 40mg twice daily, or pitavastatin 2mg or more. Low intensity statins: simvastatin 10mg, pravastatin 10-20mg, lovastatin 20mg, fluvastatin 20-40mg, pitavastatin 1mg. Statin intolerance for the purposes of this criteria is defined as documented intolerable and persistent symptoms of myalgia or signs of myositis. Two or three injections monthly will be approved for HeFH, ASCVD or Primary Hyperlipidemia. Three injections monthly will be approved for HoFH. Renewal: Chart notes, lab values documenting a decrease in LDL after initiation of therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Retevmo (selpercatinib)

Products Affected

- Retevmo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Revatio (sildenafil)

Products Affected

- Sildenafil Citrate Oral Tablet 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	Sildenafil is excluded from coverage for the treatment of Erectile Dysfunction.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Revlimid (lenalidomide)

Products Affected

- Lenalidomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Rexulti (brexpiprazole)

Products Affected

- Rexulti

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Major Depressive Disorder (MDD): Rexulti (brexpiprazole) must be used as adjunctive or add on treatment for Major Depressive Disorder (MDD), not as monotherapy. AND The patient must have a documented trial and failure of at least two (2) other formulary antidepressants for a minimum of 4 weeks. OR The patient must have a documented trial and failure of at least one (1) antidepressant and one (1) adjunctive agent, such as escitalopram, fluoxetine, paroxetine, sertraline, duloxetine, or venlafaxine for a minimum of 4 weeks, unless the patient has a documented intolerance or contraindication to the preferred medication. Schizophrenia: The patient must have a documented intolerance, side effects or lack of efficacy to at least two (2) other formulary atypical antipsychotics. Medication trials that fail due to lack of efficacy must be attempted for a minimum of 4 weeks if no response, and a minimum of 12 weeks if partial response. OR The patient has a current diagnosis of Metabolic Syndrome, Pre-Metabolic Syndrome, or Diabetes Mellitus and has failed ziprasidone or there is clinical documentation why ziprasidone is not clinically appropriate.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Rezlidhia (olutasidenib)

Products Affected

- Rezlidhia

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomsn Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Rezurock (belumosudil)

Products Affected

- Rezurock

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis of chronic graft-versus-host disease (GVHD). 2. History of failure of at least two prior lines of systemic therapy (e.g., corticosteroids, mycophenolate, tacrolimus, etc.).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	Continuation criteria: positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rinvoq (upadacitinib)

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Rheumatoid Arthritis (RA): a. DAS-28 greater than 3.2 or CDAI greater than 10.1. b. An adequate trial (3 months or more) of one of the following DMARDs: i. Hydroxychloroquine, ii. Leflunomide, iii. Methotrexate, vi. Sulfasalazine. c. Inadequate response to one or more TNF blockers (eg, Amjevita, Enbrel, Hadlima). 2. Psoriatic Arthritis (PsA): The patient has had at least a 3 month trial of ONE of the following: i. Cyclosporine, ii. Leflunomide, iii. Methotrexate. b. Inadequate response to one or more TNF blockers (eg, Amjevita, Enbrel, Hadlima). 3. Atopic Dermatitis (AD): The member is at least 12 years of age, AND b. The member has a documented diagnosis of refractory, moderate to severe AD whose disease is not adequately controlled with other systemic, and topical drug products, including: i. A medium to high potency topical steroid (e.g., mometasone, fluocinolone, fluocinonide), AND ii. A topical calcineurin inhibitor, AND c. Validated Investigators Global Assessment (vIGA-AD) score greater than or equal to 3, AND d. Eczema Area and Severity Index (EASI) score greater than or equal to 16, AND e. A minimum BSA involvement of greater than or equal to 10 percent. 4. Ulcerative Colitis (UC): a. The member has had an inadequate response to one of the following: aminosalicylates (balsalazide, mesalamine, sulfasalazine), corticosteroids, thiopurines, or cyclosporine. b. Inadequate response to at least one or more TNF blockers (e.g., Amjevita, Hadlima). 5. Ankylosing Spondylitis (AS): a. Inadequate response to an NSAID and sulfasalazine (peripheral) or NSAID alone (axial). b. The member has had an inadequate response to one or more TNF blockers(e.g., Amjevita, Enbrel, Hadlima).</p>
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist, allergist, immunologist, dermatologist, gastroenterologist

PA Criteria	Criteria Details
Coverage Duration	One (1) year
Other Criteria	6. Non-radiographic Axial Spondyloarthritis: The patient has had a documented trial and failure of a non-steroidal anti-inflammatory drug (NSAID) or such treatment is contraindicated or not tolerated. All of the following is required: 1.) Current PPD (tuberculosis) negative skin test or negative QuantiFERON-TB Gold test prior to initiation of therapy.. Continuation Criteria: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rozlytrek (entrectinib)

Products Affected

- Rozlytrek

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Rubraca (rucaparib)

Products Affected

- Rubraca

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Rydapt (midostaurin)

Products Affected

- Rydapt

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Scemblix

Products Affected

- Scemblix

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Secuado (asenapine transdermal)

Products Affected

- Secuado

PA Criteria	Criteria Details
Exclusion Criteria	Patients with dementia-related psychosis
Required Medical Information	Documentation that the patient has tried/failed two (2) formulary atypical antipsychotics.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Signifor (pasireotide)

Products Affected

- Signifor

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Patient must have a baseline 24-hour urine free cortisol (UFC) greater than 1.5 times the upper limit of normal (ULN).
Age Restrictions	18 years or older
Prescriber Restrictions	Endocrinologist
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Siklos (hydroxyurea)

Products Affected

- Siklos

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Sirturo (bedaquiline)

Products Affected

- Sirturo Oral Tablet 100 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Must be used in combination with at least 3 other drugs to which the patient's multi-drug resistant tuberculosis (MDR-TB) isolate has been shown to be susceptible in vitro.
Age Restrictions	5 years of age and weighing at least 15 kg
Prescriber Restrictions	Infectious Disease
Coverage Duration	24 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skyrizi (risankizumab-rzaa)

Products Affected

- Skyrizi (150 MG Dose)
- Skyrizi Subcutaneous
- Skyrizi Pen

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Plaque Psoriasis (PsO): a. The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). b. The disease is severe as defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. c. The patient has failed to adequately respond to, or is intolerant to a 3-month trial of a topical agent (topical corticosteroid, topical calcineurin inhibitor, topical vitamin D analogs, etc.). 2. Psoriatic Arthritis (PsA): a. An adequate trial (3 months or more) of one of the following DMARDs: i. Cyclosporine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine 3. Crohn's Disease (CD): a. For induction and maintaining clinical remission in patients with moderately to severely active Crohns Disease who have had an inadequate response or intolerance to conventional therapy. For the purpose of this policy, conventional therapy includes the use of ONE of the following: i. Corticosteroids (e.g., prednisone, prednisolone, dexamethasone, budesonide). ii. Methotrexate iii. Thiopurines (azathioprine, mercaptopurine) b. Documentation that the member has received approval through their Part B benefit for the office-administered induction treatment must be received before PHP can approve the self-administered maintenance treatment.</p>
Age Restrictions	
Prescriber Restrictions	Dermatologist, gastroenterologist, rheumatologist
Coverage Duration	1 year

PA Criteria	Criteria Details
Other Criteria	For all Indications: 1) Current PPD (tuberculosis) negative skin test, negative QuantiFERON-TB Gold test, or documented treatment for latent tuberculosis prior to initiation of therapy. Continuation of Therapy Criteria: Documentation of clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Solaraze

Products Affected

- Diclofenac Sodium External Gel 3 %
- Diclofenac Sodium Transdermal Gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of osteoarthritis, pain
Required Medical Information	Documented diagnosis of actinic keratosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	90 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Somavert (pegvisomant)

Products Affected

- Somavert

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of acromgaly by 1) serum growth hormone (GH) level greater than 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis, or 2) elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician) at the time of diagnosis. The member must have also had an inadequate response to surgery, radiotherapy, or a dopamine agonist (e.g., cabergoline, bromocriptine), or is not a candidate for surgery, radiotherapy, or dopamine agonist therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 3 months Continuation: 1 year
Other Criteria	Continuation criteria: Documentation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sprycel (dasatinib)

Products Affected

- Sprycel

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Stelara (ustekinumab)

Products Affected

- Stelara Intravenous
- Stelara Subcutaneous Solution 45 MG/0.5ML
- Stelara Subcutaneous Solution Prefilled Syringe

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of moderate to severe Plaque Psoriasis (PsO). The patient must have greater than 3% of their body surface area (BSA) affected by plaque psoriasis). The disease is defined by a total Psoriasis Area Severity Index (PASI) of at least 5 or more and/or a Dermatology Life Quality Index (DLQI) of more than 5. The patient has failed to adequately respond to two of the following: Amjevita, Cosentyx, Enbrel, Hadlima, or Skyrizi.</p> <p>Diagnosis of active Psoriatic Arthritis (PsA). The patient has failed to adequately respond to two of the following: Amjevita, Cosentyx, Enbrel, Hadlima, Rinvoq (must have tried a TNF-inhibitor), Skyrizi, or Xeljanz (must have tried a TNF-inhibitor).</p> <p>Crohn's Disease (CD): For induction and maintaining clinical remission in patients with moderately to severely active Crohns Disease who have had an inadequate response or intolerance to conventional therapy. For the purpose of this policy, conventional therapy includes the use of ONE of the following: i. Corticosteroids (e.g., prednisone, prednisolone, dexamethasone, budesonide). ii. Methotrexate iii. Thiopurines (azathioprine, mercaptopurine). Inadequate response to Amjevita, Skyrizi.</p> <p>Diagnosis of moderate to severe active Ulcerative Colitis (UC). The member has had a failed to have an adequate response to two of the following: Amjevita, Hadlima and Rinvoq (must have tried a TNF-inhibitor) or Xeljanz (must have tried a TNF-inhibitor).</p>
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a Dermatologist, a Rheumatologist or a Gastroenterologist

PA Criteria	Criteria Details
Coverage Duration	1 yr
Other Criteria	Documentation of a positive clinical response to the approved therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Stivarga (regorafenib)

Products Affected

- Stivarga

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Strensiq (asfotase alfa)

Products Affected

- Strensiq Subcutaneous Solution 40 MG/ML, 80 MG/0.8ML

PA Criteria	Criteria Details
Exclusion Criteria	Adult-onset hypophosphatasia
Required Medical Information	Documented diagnosis of perinatal/infantile onset or juvenile-onset hypophosphatasia (HPP) confirmed by all of the following: a)Patient is less than or equal to 18 at age of onset of disease, b)Radiographic evidence of HPP, c)Rachitic deformities, d)premature loss of primary teeth prior to age 5, e)Delay in skeletal growth resulting in delay of motor development, f)History or presence of non-traumatic fractures or delayed fracture healing. Molecular genetic testing and results for mutation(s) in the ALPL gene. Baseline serum alkaline phosphatase (ALP), below normal range for patients age. Serum vitamin B-6 levels, elevated and the patient has not received vitamin B6 supplementation in the previous week. Elevated serum or urine Phosphoethanolamine (PEA) level. Baseline ophthalmologic exam and renal ultrasound being monitored for signs and symptoms of ectopic calcifications.
Age Restrictions	
Prescriber Restrictions	Must be prescribed by, or in consultation with, an endocrinologist, geneticist or a metabolic specialist.
Coverage Duration	6 months
Other Criteria	Continuation criteria: Documentation that the patient is tolerating treatment and responding to treatment, as evidenced by improvement in respiratory status, or radiographic findings. Dose is not to exceed FDA label maximum. Strensiq can be injected three times per week or six times per week. Strensiq is only covered as a three times per week injection.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Sutent (sunitinib)

Products Affected

- SUNItinib Malate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Sylatron (peginterferon alfa-2b)

Products Affected

- Sylatron Subcutaneous Kit 200 MCG, 300 MCG, 600 MCG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or there must be a Class I or IIa recommendation in the Thomson Micromedex DrugDex compendium.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Symdeko (ivacaftor / tezacaftor)

Products Affected

- Symdeko

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	6 years of age and older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Synarel (nafarelin acetate)

Products Affected

- Synarel

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Endometriosis: Documented contraindication, intolerance, or treatment failure with Lupron Depot (leuprolide).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tabloid (thioguanine)

Products Affected

- Tabloid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tabrecta (capmatinib)

Products Affected

- Tabrecta

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tafinlar (dabrafenib)

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tagrisso (osimertinib)

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Takhzyro (lanadelumab-flyo)

Products Affected

- Takhzyro

PA Criteria	Criteria Details
Exclusion Criteria	Not to be used in combination with other prophylactics (Cinryze or Haegarda)
Required Medical Information	A diagnosis of hereditary angioedema (HAE) has been clinically established by or in consultation with an Allergist or Immunologist.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	4 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Talzenna (talazoparib)

Products Affected

- Talzenna

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tarceva (erlotinib)

Products Affected

- Erlotinib HCl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Targretin (bexarotene)

Products Affected

- Bexarotene

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tasigna (nilotinib)

Products Affected

- Tasigna

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tazverik (tazemetostat)

Products Affected

- Tazverik

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	must be at least 16 years old
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tepmetko (tepotinib)

Products Affected

- Tepmetko

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Testosterone injection

Products Affected

- Testosterone Cypionate Intramuscular Solution 100 MG/ML, 200 MG/ML, 200 MG/ML (1 ML)
- Testosterone Enanthate Intramuscular Solution

PA Criteria	Criteria Details
Exclusion Criteria	Testosterone replacement will not be covered for the treatment of sexual dysfunction.
Required Medical Information	If using for primary hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with elevated luteinizing hormone (LH) and follicular stimulating hormone (FSH) levels. If using for hypogonadotropic hypogonadism, the patient must have two low total testosterone levels (drawn on separate days) with low to low-normal LH and FSH levels.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Thalomid (thalidomide)

Products Affected

- Thalomid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tibsovo (ivosidenib)

Products Affected

- Tibsovo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Truqap (capivasertib)

Products Affected

- Truqap

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Six (6) months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Truseltiq (infigratinib)

Products Affected

- Truseltiq (100MG Daily Dose)
- Truseltiq (125MG Daily Dose)
- Truseltiq (50MG Daily Dose)
- Truseltiq (75MG Daily Dose)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tukysa (tucatinib)

Products Affected

- Tukysa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Turalio (pexidartinib)

Products Affected

- Turalio Oral Capsule 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tykerb (lapatinib ditosylate)

Products Affected

- Lapatinib Ditosylate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tymlos (abaloparatide)

Products Affected

- Tymlos

PA Criteria	Criteria Details
Exclusion Criteria	Cumulative use of Parathyroid Hormone Analogs greater than 2 years will not be approved.
Required Medical Information	A diagnosis of osteoporosis and a T-score of -2.5 or less at the femoral neck, total hip or lumbar spine by DXA. The patient has failed or is intolerant to other available osteoporosis therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Uceris (budesonide tablets)

Products Affected

- Budesonide ER Oral Tablet Extended Release 24 Hour

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. The member has a diagnosis of active mild to moderate ulcerative colitis, AND 2. The member must have had previous treatment or intolerance to at least two of the following: sulfasalazine, balsalazide, or mesalamine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	8 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ukoniq (umbralisib)

Products Affected

- Ukoniq

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or there must be a Class I or IIa recommendation in the Thomson Micromedex DrugDex compendium.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Uloric (febuxostat)

Products Affected

- Febuxostat

PA Criteria	Criteria Details
Exclusion Criteria	Current azathioprine or mercaptopurine use.
Required Medical Information	1. Documented diagnosis of gout. 2. One of the following must be met: a. Documented failure at maximally tolerated dose of allopurinol - a documented failure is considered as non-resolution of tophi or at least 4 gout attacks (joint flares) per year - OR - b. documented intolerance to allopurinol (examples of intolerance include skin reactions or cytopenias).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Uptravi (selexipag)

Products Affected

- Uptravi Oral
- Uptravi Titration

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Initial: Diagnosis of WHO Group I pulmonary arterial hypertension (PAH) AND Patient is symptomatic. a) Diagnosis of pulmonary arterial hypertension was confirmed by right heart catheterization OR b) patient is currently on any therapy for the diagnosis of pulmonary arterial hypertension. Reauth: Documentation of positive clinical response to Uptravi therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Reauthorization: 1 year
Other Criteria	Initial: One of the following: a) History of inadequate response, contraindication, or intolerance to a formulary PDE5 inhibitor, and history of inadequate response, contraindication, or intolerance to a formulary endothelin receptor antagonist b) For continuation of prior Uptravi therapy. Initial/Reauth: Not taken in combination with a prostanoid/prostacyclin analogue (eg, eprostenol, iloprost, treprostinil).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Valchlor (mechlorethamine)

Products Affected

- Valchlor

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Vanflyta (quizartinib)

Products Affected

- Vanflyta

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium, or one of the following: a Class I, IIA, or IIB recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Venclexta (venetoclax)

Products Affected

- Venclexta
- Venclexta Starting Pack

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Verquvo (vericiguat)

Products Affected

- Verquvo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis heart failure classified as one of the following: a. New York Heart Association Class II, III, or IV. 2. Ejection fraction is less than 45 percent. 3. One of the following: a. hospitalization for heart failure within the past six months. b. outpatient IV diuretics for heart failure within the past three (3) months. 4. patient is on a stabilized dose and receiving concomitant therapy with a maximally tolerated dose of all of the following: (unless contraindicated or not tolerated): a. beta-blocker (eg., bisoprolol, carvedilol, metoprolol) b. angiotensin converting enzyme (ACE) inhibitor (e.g., captopril, lisinopril), angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan), or angiotensin receptor-neprilysin inhibitor (ARNI) (e.g., Entresto), c. aldosterone antagonist (e.g., spironolactone).
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist.
Coverage Duration	One (1) year
Other Criteria	Continuation criteria: positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verzenio (abemaciclib)

Products Affected

- Verzenio

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Vfend (voriconazole)

Products Affected

- Voriconazole Intravenous
- Voriconazole Oral Tablet 200 MG
- Voriconazole Oral Suspension Reconstituted

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of invasive aspergillosis, candidemia, esophageal candidiasis, a disseminated (widespread) Candida infection in the skin, or a Candida infection in the abdomen, kidney, bladder wall, or wounds and a documented trial/failure of fluconazole. Documented fungal infection caused by Fusariosis or Scedosporium species. Culture and sensitivity report.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vitrakvi (larotrectinib)

Products Affected

- Vitrakvi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Vizimpro (dacomitinib)

Products Affected

- Vizimpro

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Vonjos (pacritinib)

Products Affected

- Vonjo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Votrient (pazopanib)

Products Affected

- PAZOPanib HCl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Vraylar (cariprazine)

Products Affected

- Vraylar

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	The patient must have a documented intolerance, side effects or lack of efficacy to at least two (2) other formulary atypical antipsychotics. Medication trials that fail due to lack of efficacy must be attempted for a minimum of 4 weeks if no response, and a minimum of 12 weeks if partial response, unless the patient has a documented intolerance or contraindication to the preferred medication. OR The patient has a current diagnosis of Metabolic Syndrome, Pre-Metabolic Syndrome, or Diabetes Mellitus and has failed ziprasidone or there is clinical documentation why ziprasidone is not clinically appropriate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vyndaqel (tafamidis meglumine)

Products Affected

- Vyndaqel

PA Criteria	Criteria Details
Exclusion Criteria	Less than 18 years of age, concomitant use with patisiran or inotersen.
Required Medical Information	For the treatment of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis confirmed by biopsy and DNA mutation analysis and the patient has a medical history of heart failure with at least one hospitalization for heart failure. New York Heart Association (NYHA) class I, II or III heart failure symptoms.
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or a physician who specializes in the treatment of amyloidosis.
Coverage Duration	Initial: 6 months Renew: 1 year
Other Criteria	Renew: Patient has experienced a positive clinical response (e.g. cardiac function, serum TTR levels).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vyvanse (lisdexamphetamine)

Products Affected

- Lisdexamphetamine Dimesylate Oral
Capsule 10 MG, 20 MG, 30 MG, 40 MG,
50 MG, 60 MG, 70 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Attention Deficit Hyperactivity Disorder (ADHD): a. diagnosis of ADHD. b. Patient has had an inadequate response to at least one-month trial of at least two of the following generic medications: amphetamine, amphetamine/dextroamphetamine, dexamethylphenidate, dextroamphetamine, methylphenidate. 2. Binge Eating Disorder (BED): a. diagnosis of BED. b. Inadequate response to a selective serotonin reuptake inhibitor (SSRI) or topiramate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	One (1) year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Welireg (belzurifan)

Products Affected

- Welireg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Xalkori (crizotinib)

Products Affected

- Xalkori

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Xatmep (methotrexate)

Products Affected

- Xatmep

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting a trial/failure or contraindication to methotrexate tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xeljanz (tofacitinib)

Products Affected

- Xeljanz
- Xeljanz XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. Rheumatoid Arthritis (RA): a. Documented presence of moderate to severe RA: DAS-28 greater than 3.2 or CDAI greater than 10.1. b. An adequate trial (3 months or more) of one of the following DMARDs: i. Hydroxychloroquine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine c. Trial and failure, unless contraindicated or not tolerated, to at least one tumor necrosis factor (TNF) blocker (e.g., Amjevita, Enbrel, Hadlima). 2. Psoriatic Arthritis (PsA): a. An adequate trial (3 months or more) of one of the following DMARDs: i. Cyclosporine ii. Leflunomide iii. Methotrexate iv. Sulfasalazine b. Trial and failure, unless contraindicated or not tolerated, to at least one tumor necrosis factor (TNF) blocker (e.g., Amjevita, Enbrel, Hadlima). 3. Ulcerative Colitis (UC): a. The patient must have an adequate trial (3 months or more) or intolerance to ONE of the following: i. 5-Aminosalicylates (balsalazide, mesalamine, sulfasalazine) ii. Cyclosporine iii. Steroids iv. Thiopurines (azathioprine, 6-MP) b. Trial and failure to at least one tumor necrosis factor (TNF) blocker (e.g., Amjevita, Hadlima). 4. Juvenile Idiopathic Arthritis (JIA): a. An adequate trial (3 months or more) of one of the following DMARDs: i. Leflunomide ii. Methotrexate iii. Sulfasalazine b. Trial and failure to at least one tumor necrosis factor (TNF) blocker (e.g., Amjevita, Enbrel, Hadlima). 5. Ankylosing Spondylitis (AS): a. The patient has a documented trial and failure with a non-steroidal anti-inflammatory drug (NSAID) or such treatment is contraindicated or not tolerated. b. Patients with peripheral arthritis must have a documented trial and failure with sulfasalazine or such treatment is contraindicated or not tolerated. c. Patients with axial disease and a trial a trial/failure of NSAIDs can be started on Xeljanz. d. 5. Trial and failure of at least one tumor necrosis factor (TNF) blocker (e.g., Amjevita, Enbrel, Hadlima).</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	RA//JIA/AS- prescribed by or in consultation with a rheumatologist, PsA- prescribed by or in consultation with a dermatologist or rheumatologist, UC- prescribed by or in consultation with a gastroenterologist
Coverage Duration	One (1) year
Other Criteria	For all Indications: 1) Current PPD (tuberculosis) negative skin test, negative QuantiFERON-TB Gold test, or documented treatment for latent tuberculosis prior to initiation of therapy. Continuation of Therapy Criteria: Documentation of clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xenazine (tetrabenazine)

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Exclusion Criteria	Tetrabenazine is not covered for patients who are actively suicidal, who have untreated or inadequately treated depression, who have impaired hepatic function, or who are currently taking monoamine oxidase inhibitors or reserpine.
Required Medical Information	Treatment of chorea associated with Huntington's disease. Documentation that member is being monitored for depression and suicidal ideation. Renewal: Chart notes documenting that the patient's disease has improved based on prescriber's assessment while on therapy.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 3 months Renewal: 1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xermelo (telotristat)

Products Affected

- Xermelo

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chart notes documenting ALL of the following: 1)A diagnosis of carcinoid syndrome diarrhea, 2)An inadequate treatment response to a somatostatin analog (SSA), at a maximum tolerated dose, after at least 3 months of therapy, 3)A documented trial/failure of adjunct treatment with an anti-diarrheal medication, such as loperamide or ondansetron, 4)Chart notes documenting at least four bowel movements per day, 5)Xermelo will be used by the patient in combination with a SSA.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Oncologist or Gastroenterologist
Coverage Duration	Initial: 12 weeks Renewal: 1 year
Other Criteria	Renewal: Diagnosis of carcinoid syndrome diarrhea, and the patient will continue to use Xermelo with a SSA. Chart notes documenting a decrease in baseline in the amount of daily bowel movements.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xgeva (denosumab)

Products Affected

- Xgeva

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xifaxan (rifaximin)

Products Affected

- Xifaxan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	If using for traveler's diarrhea (TD), trial and failure of an oral antibiotic such as Zithromax (azithromycin), Levaquin (levofloxacin) or Floxin (ofloxacin). If using for hepatic encephalopathy (HE), trial and failure or documented intolerance/contraindication to lactulose oral solution. If using for Irritable Bowel Syndrome, Diarrhea Predominant (IBS-D) documented diagnosis of IBS-D, trial and failure of at least two (2) of the following: anti-diarrheals, antispasmodics, or tricyclic antidepressants.
Age Restrictions	TD: 12 years or older HE: 18 years or older IBS-D: 18 years or older
Prescriber Restrictions	
Coverage Duration	TD: 3 days per request HE: 1 year IBS-D: 14 days per request
Other Criteria	Quantity limits will be diagnosis dependent: TD three times daily, HE twice daily dosing, IBS-D three times daily.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xiidra (lifitegrast)

Products Affected

- Xiidra

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of dry eye disease and the patient has suppressed tear production due to ocular inflammation as determined by at least one of the following diagnostic tests: Schirmer Tear Test (STT), Corneal Fluorescein Staining (CFS), tear break-up time, tear film osmolarity, ocular surface dye staining.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months Renewal: 1 year
Other Criteria	Renewal: Documentation of positive clinical response to Xiidra therapy (increased tear production or improvement in dry eye symptoms).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xolair (omalizumab)

Products Affected

- Xolair

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	If using for persistent asthma (PA), medical records must document all of the following: IgE level greater than or equal to 30 IU/mL, specific evidence of allergic asthma, supported by clinical and lab findings such as positive skin tests, symptom patterns, etc., inadequate control with an inhaled corticosteroid and a long acting beta-2 agonist combination therapy, and evidence of persistent symptoms requiring frequent rescue therapy, practitioner visits despite inhaled corticosteroids, ER visits OR inadequate control OR intolerance OR contraindication to inhaled corticosteroid and a long acting beta-2 agonist combination. If using for chronic idiopathic urticaria (CIU) (initial request): Documentation of the following: 30-day trial of a second generation non-sedating anti-histamine and one of the following: a leukotriene antagonist or a Histamine-2 (H2) antagonist, OR intolerance OR contraindication to preferred medications. CIU (renewal request): Documentation of a reduction in exacerbation frequency and intensity. For nasal polyps, must have bilateral polyps as determined by a nasal polyp score (NPS) greater than or equal to 5 with NPS greater than or equal to 2 in each nostril, and a documented inadequate response to nasal corticosteroids.
Age Restrictions	
Prescriber Restrictions	Allergist, Pulmonologist, Dermatologist, Immunologist, or Otolaryngologist.
Coverage Duration	Initial: six (6) months. Renewal: one (1) year.
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Xospata (gilteritinib)

Products Affected

- Xospata

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Xpovio (selinexor)

Products Affected

- Xpovio (100 MG Once Weekly)
- Xpovio (40 MG Once Weekly)
- Xpovio (40 MG Twice Weekly)
- Xpovio (60 MG Once Weekly)
- Xpovio (60 MG Twice Weekly)
- Xpovio (80 MG Once Weekly)
- Xpovio (80 MG Twice Weekly)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Xtandi (enzalutamide)

Products Affected

- Xtandi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1. Diagnosis of metastatic, castration-resistant prostate cancer AND history of failure, contraindication, or intolerance to abiraterone (Zytiga), OR diagnosis of metastatic, castration-sensitive prostate cancer AND history of failure, contraindication, or intolerance to abiraterone (Zytiga), OR diagnosis of non-metastatic, castration-resistant prostate cancer AND history of failure, contraindication, or intolerance to Nubeqa.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Xyrem (sodium oxybate)

Products Affected

- Xyrem

PA Criteria	Criteria Details
Exclusion Criteria	Will not be approved in combination with sedative hypnotics or alcohol, or if patient has a succinic semialdehyde dehydrogenase deficiency.
Required Medical Information	Adults: If using for excessive daytime sleepiness in narcolepsy, trial and failure of, or intolerance to a cerebral stimulant (methylphenidate or dextroamphetamine) and Nuvigil (armodafinil). Children will not be required to try armodafinil.
Age Restrictions	7 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Yonsa (abiraterone)

Products Affected

- Yonsa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zejula (niraparib)

Products Affected

- Zejula

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zelboraf (vemurafenib)

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zolinza (vorinostat)

Products Affected

- Zolinza

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zonisade(zonisamide)

Products Affected

- Zonisade

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	The member must have tried and failed two formulary anti-convulsants, with one agent being the oral capsule dosage form of zonisamide, unless contraindicated or not tolerated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	Some FDA-approved Indications Only.
Off Label Uses	
Part B Prerequisite	No

Ztalmy (ganaxolone)

Products Affected

- Ztalmy

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD). Patient has a mutation in the CDKL5 gene. Documentation of baseline monthly seizure frequency. Trial and failure, contraindication, or intolerance to two formulary anticonvulsants (e.g., valproic acid, levetiracetam, lamotrigine).
Age Restrictions	Patient is 2 years of age or older.
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	One (1) year
Other Criteria	Continuation: Documentation of a sustained reduction in monthly seizure frequency compared to baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zurzuvae (zuranolone)

Products Affected

- Zurzuvae

PA Criteria	Criteria Details
Exclusion Criteria	Current pregnancy, past medical history of bipolar disorder, schizophrenia, or schizoaffective disorder.
Required Medical Information	Documentation of moderate to severe postpartum depression diagnosis and submission of validated screening tool (e.g., EPDS, PHQ-9). Documentation that member has not had a major depressive episode prior to third trimester of pregnancy and not later than the first 4 weeks following delivery. Member has tried and failed a formulary SSRI or SNRI for postpartum depression.
Age Restrictions	18 years of age
Prescriber Restrictions	Prescribed by or in consultation with a psychiatrist or OB/GYN.
Coverage Duration	One (1) month
Other Criteria	Reauthorization: Only one course of 14 days will be allowed per 12 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zydelig (idelalisib)

Products Affected

- Zydelig

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zykadia (ceritinib)

Products Affected

- Zykadia Oral Tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Non-FDA approved indications, there must be a Category 1 or 2A recommendation in the National Comprehensive Cancer Network (NCCN) compendium or, one of the following: a Class I, Iia, or Iib recommendation in the Thomson Micromedex DrugDex compendium, narrative text that is supportive by the AFHS-DI or Clinical Pharmacology, or an Evidence Level A by Lexi-Drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Zyvox (linezolid)

Products Affected

- Linezolid Oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical records documenting all of the following: Prescribed by or in consultation with an infectious disease specialist. Culture and sensitivity report documents that the isolated pathogen is susceptible to linezolid and the patient has failed one formulary antibiotic to which the isolated pathogen is susceptible per the culture and sensitivity report. OR the request is for continuation of therapy that was initiated in an acute care hospital from which the patient was discharged.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Non-VREF: 14 days VREF: 28 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Index

A

Abilify MyCite..... 2

Abilify MyCite Maintenance Kit Oral Tablet
Therapy Pack 15 MG, 2 MG, 20 MG, 5
MG 2

Abilify MyCite Starter Kit Oral Tablet
Therapy Pack 10 MG 2

Actemra ACTPen..... 4

Actemra Subcutaneous..... 4

Adempas 7

Aimovig 38

Akeega 10

Alecensa 11

Alora 96, 97

Alosetron HCl 126

Alunbrig 12

Alyq..... 6

Amabelz 96, 97

Ambrisentan 169

Amitriptyline HCl Oral 93

Amjevita Subcutaneous Solution Auto-
Injector 13, 14

Amjevita Subcutaneous Solution Prefilled
Syringe 40 MG/0.4ML..... 13, 14

Amjevita-Ped 10kg to <15kg 13, 14

Amjevita-Ped 15kg to <30kg 13, 14

Amoxapine 93

Androderm Transdermal Patch 24 Hour... 15

Apokyn Subcutaneous Solution Cartridge 17

Apomorphine HCl Subcutaneous 17

Aprepitant Oral Capsule 59

Arcalyst..... 18

ARIPiprazole Oral Tablet Dispersible 3

Armodafinil..... 147

Augtyro 20

Austedo 21

Austedo Patient Titration Kit 21

Austedo XR..... 21

Austedo XR Patient Titration..... 21

Auvelity..... 22

Ayvakit..... 23

B

Balversa..... 24

Benlysta Subcutaneous 25

Besremi 26

Bexarotene 215

Bosentan..... 169

Bosulif..... 27

Braftovi 28

Bronchitol 29

Brukinsa 30

Budesonide ER Oral Tablet Extended
Release 24 Hour 228

Budesonide Oral.....	64
C	
Cabometyx	32
Calquence.....	33
Caplyta	34
Caprelsa.....	35
Carglumic Acid.....	36
Cayston	37
chlordiazepoxide HCl.....	95
chlorpromazine HCl Oral Tablet	95
Cinryze.....	39
clomipramine HCl Oral.....	93
CombiPatch.....	96, 97
Cometriq (100 MG Daily Dose)	43
Cometriq (140 MG Daily Dose)	43
Cometriq (60 MG Daily Dose)	43
Copiktra.....	44
Corlanor	45
Cosentyx (300 MG Dose)	46, 47
Cosentyx Sensoready (300 MG).....	46, 47
Cosentyx Sensoready Pen.....	46, 47
Cosentyx Subcutaneous	46, 47
Cosentyx UnoReady	46, 47
Cotellic.....	48
Cyclobenzaprine HCl Oral.....	79
Cystaran	49

D	
Daurismo.....	52
Deferasirox Oral Tablet	53
Deferasirox Oral Tablet Soluble	53
Deferiprone	76
Desipramine HCl Oral	93
Diclofenac Sodium External Gel 3 %	196
Diclofenac Sodium Transdermal Gel 3 %	196
Dificid Oral Tablet	55
Dotti	96, 97
Doxepin HCl Oral Capsule	93
Drizalma Sprinkle	56
Dronabinol	132
Droxidopa	142
Dupixent.....	57, 58
E	
Emend Oral Suspension Reconstituted.....	59
Emgality	60
Enbrel Mini	61, 62
Enbrel Subcutaneous Solution 25 MG/0.5ML	61, 62
Enbrel Subcutaneous Solution Prefilled Syringe	61, 62
Enbrel Subcutaneous Solution Reconstituted	61, 62
Enbrel SureClick Subcutaneous Solution Auto-Injector.....	61, 62

Endari	63	Flebogamma DIF	83
Epidiolex	66	Forteo Subcutaneous Solution 600 MCG/2.4ML	80
Erivedge	67	Forteo Subcutaneous Solution Pen-Injector	80
Erleada	68	Fotivda	81
Erlotinib HCl.....	214	Fruzaqla.....	82
Estradiol Oral Tablet 2 MG	96, 97	Fyavolv	96, 97
Estradiol-Norethindrone Acet.....	96, 97	G	
Estropipate Oral	96, 97	Gamunex-C	83
Evamist	96, 97	Gattex	84
Everolimus Oral Tablet 10 MG, 2.5 MG, 5 MG, 7.5 MG.....	8	Gavreto.....	85
Everolimus Oral Tablet Soluble.....	9	Gefitinib	106
Exkivity	71	Gilotrif.....	86
Extavia Subcutaneous Kit	72	H	
F		Hadlima.....	89, 90
Farydak	73	Hadlima PushTouch.....	89, 90
Fasenra Pen	74	Hexalen	92
Fasenra Subcutaneous Solution Prefilled Syringe 30 MG/ML.....	74	I	
Febuxostat	230	Ibrance.....	98
Fensolvi.....	75	Icatibant Acetate	77
Fensolvi (6 Month)	75	Iclusig.....	99
fentaNYL Citrate Buccal Lozenge On A Handle 1200 MCG, 400 MCG, 600 MCG, 800 MCG	5	IDHIFA	100
FentaNYL Citrate Buccal Lozenge On A Handle 1600 MCG	5	Imatinib Mesylate	87
Firdapse.....	78	Imbruvica Oral Capsule	101
		Imbruvica Oral Suspension.....	101
		Imbruvica Oral Tablet 140 MG, 280 MG, 420 MG	101

Imipramine HCl Oral	93
Increlex	102
Inlyta	103
Inqovi	104
Inrebic	105
Ivermectin Oral	107
Iwilfin.....	108

J

Jakafi.....	109
Jaypirca	110
Jevantique Lo.....	96, 97
Jinteli.....	96, 97
Juxtapid.....	111
Jylamvo.....	112

K

Kalydeco Oral Packet 25 MG, 50 MG, 75 MG	113
Kalydeco Oral Tablet.....	113
Kerendia.....	114
Kineret Subcutaneous Solution Prefilled Syringe	115
Kisqali (200 MG Dose).....	116
Kisqali (400 MG Dose).....	116
Kisqali (600 MG Dose).....	116
Kisqali 200 Dose.....	116
Kisqali 400 Dose.....	116
Kisqali 600 Dose.....	116

Kisqali Femara (200 MG Dose).....	116
Kisqali Femara (400 MG Dose).....	116
Kisqali Femara (600 MG Dose).....	116
Koselugo	118
Krazati.....	119

L

Lapatinib Ditosylate.....	226
Lenalidomide	179
Lenvima (10 MG Daily Dose)	121
Lenvima (12 MG Daily Dose)	121
Lenvima (14 MG Daily Dose)	121
Lenvima (18 MG Daily Dose)	121
Lenvima (20 MG Daily Dose)	121
Lenvima (24 MG Daily Dose)	121
Lenvima (4 MG Daily Dose)	121
Lenvima (8 MG Daily Dose)	121
Leukine Injection Solution Reconstituted	122
Leukine Intravenous.....	122
Lidocaine External Patch 5 %	123
Linezolid Oral	270
Lisdexamfetamine Dimesylate Oral Capsule 10 MG, 20 MG, 30 MG, 40 MG, 50 MG, 60 MG, 70 MG.....	244
Lonsurf.....	124
Lopreeza.....	96, 97
Lorbrena.....	125
Lumakras.....	127

Lybalvi	128
Lyllana	96, 97
Lynparza	129
Lytgobi (12 MG Daily Dose).....	131
Lytgobi (16 MG Daily Dose).....	131
Lytgobi (20 MG Daily Dose).....	131

M

Matulane	133
Mavyret	134
Mekinist	135
Mektovi	136
Menest.....	96, 97
miFEPRIStone Oral Tablet 300 MG.....	117
Mimvey	96, 97
Mimvey Lo.....	96, 97
Mounjaro.....	88
Movantik.....	137

N

Natpara	138
Nerlynx	139
Ninlaro	141
Nitisinone.....	158
Norethindrone-Eth Estradiol.....	96, 97
Nortriptyline HCl Oral Capsule	93
Nortriptyline HCl Oral Solution	93
Nubeqa	144
Nuedexta	145

Y0055_MPC092232_NSR_C_09232022
Formulary ID 00024541
Version 021
Last Updated: 06/20/2024Page | 275

Nuplazid Oral Capsule	146
Nuplazid Oral Tablet 10 MG	146

O

Ocrevus	148
Odomzo.....	149
Ofev.....	150
Ogsiveo Oral Tablet 50 MG	151
Ojjaara.....	152
Omnitrope	153, 154
Onureg.....	155
Opsumit.....	156
Orencia ClickJect	157
Orencia Subcutaneous Solution Prefilled Syringe	157
Orgovyx	159
Orkambi Oral Tablet.....	160
Orserdu.....	161
Otezla	162
Ozempic (0.25 or 0.5 MG/DOSE)	88
Ozempic (1 MG/DOSE) Subcutaneous Solution Pen-Injector 4 MG/3ML.....	88
Ozempic (2 MG/DOSE)	88

P

PARoxetine HCl	94
PARoxetine HCl ER	94
Paxil Oral Suspension.....	94
PAZOPanib HCl	241

Pemazyre.....	163	Repatha	175, 176
penicillAMINE Oral Tablet	54	Repatha Pushtonex System.....	175, 176
Perseris	164	Repatha SureClick	175, 176
Piqray (200 MG Daily Dose).....	165	Retacrit Injection Solution 10000	
Piqray (250 MG Daily Dose).....	165	UNIT/ML, 10000 UNIT/ML(1ML), 2000	
Piqray (300 MG Daily Dose).....	165	UNIT/ML, 3000 UNIT/ML, 4000	
Pirfenidone Oral Capsule	70	UNIT/ML, 40000 UNIT/ML	69
Pirfenidone Oral Tablet 267 MG, 801 MG70		Retevmo	177
Pomalyst.....	166	Rexulti.....	180, 181
Posaconazole Oral.....	143	Rezlidhia	182
Prefest	96, 97	Rezurock	183
Pregabalin ER	130	Rinvoq.....	184, 185
Premarin Oral.....	96, 97	Roflumilast.....	50
Premphase	96, 97	Rozlytrek.....	186
Prempro.....	96, 97	Rubraca	187
Prolastin-C	167	Rybelsus	88
Promacta	168	Rydapt.....	188
Protriptyline HCl.....	93	S	
Purixan	170	Sajazir	77
Pyrimethamine Oral	51	Sapropterin Dihydrochloride	120
Q		Scemblix	189
Qinlock.....	171	Scopolamine.....	95
QuiNINE Sulfate Oral.....	172	Secuado	190
R		Signifor	191
Regranex	173	Siklos.....	192
Relistor Oral.....	174	Sildenafil Citrate Oral Tablet 20 MG	178
Relistor Subcutaneous Solution	174	Sirturo Oral Tablet 100 MG.....	193

Skyrizi Pen	194, 195	Takhzyro	212
Skyrizi Subcutaneous.....	194, 195	Talzenna.....	213
Sodium Phenylbutyrate Oral Powder 3 GM/TSP	31	Tasigna.....	216
Sodium Phenylbutyrate Oral Tablet.....	31	Tasimelteon.....	91
Sofosbuvir-Velpatasvir	65	Tazverik	217
Somavert	197	Tepmetko	218
SORafenib Tosylate.....	140	Teriflunomide	19
Sprycel	198	Teriparatide (Recombinant) Subcutaneous Solution Pen-Injector 620 MCG/2.48ML	80
Stelara Intravenous.....	199, 200	Testosterone Cypionate Intramuscular Solution 100 MG/ML, 200 MG/ML, 200 MG/ML (1 ML)	219
Stelara Subcutaneous Solution 45 MG/0.5ML.....	199, 200	Testosterone Enanthate Intramuscular Solution.....	219
Stelara Subcutaneous Solution Prefilled Syringe	199, 200	Testosterone Transdermal Gel 1.62 %, 10 MG/ACT (2%), 12.5 MG/ACT (1%), 20.25 MG/1.25GM (1.62%), 20.25 MG/ACT (1.62%), 25 MG/2.5GM (1%), 40.5 MG/2.5GM (1.62%), 50 MG/5GM (1%).....	16
Stivarga	201	Tetrabenazine	250
Strensiq Subcutaneous Solution 40 MG/ML, 80 MG/0.8ML	202, 203	Thalomid.....	220
SUNItinib Malate.....	204	Tibsovo	221
Sylatron Subcutaneous Kit 200 MCG, 300 MCG, 600 MCG	205	Trientine HCl	40
Symdeko	206	Trimipramine Maleate Oral	93
Synarel	207	Trulicity.....	88
T		Truqap.....	222
Tabloid	208	Truseltiq (100MG Daily Dose).....	223
Tabrecta.....	209	Truseltiq (125MG Daily Dose).....	223
Tadalafil (PAH)	6		
Tafinlar.....	210		
Tagrisso.....	211		

Truseltiq (50MG Daily Dose).....	223
Truseltiq (75MG Daily Dose).....	223
Tukysa.....	224
Turalio Oral Capsule 125 MG	225
Tymlos	227
U	
Udenyca Subcutaneous Solution Prefilled Syringe	41, 42
Ukoniq.....	229
Uptravi Oral	231
Uptravi Titration	231
V	
Valchlor.....	232
Vanflyta.....	233
Venclexta	234
Venclexta Starting Pack	234
Verquvo.....	235
Verzenio	236
Victoza Subcutaneous Solution Pen-Injector	88
Vitrakvi	238
Vizimpro	239
Vonjo.....	240
Voriconazole Intravenous	237
Voriconazole Oral Suspension Reconstituted	237
Voriconazole Oral Tablet 200 MG	237
Vraylar	242

Vyndaqel.....	243
W	
Welireg.....	245
X	
Xalkori	246
Xatmep.....	247
Xeljanz	248, 249
Xeljanz XR.....	248, 249
Xermelo.....	251
Xgeva	252
Xifaxan.....	253
Xiidra	254
Xolair	255, 256
Xospata	257
Xpovio (100 MG Once Weekly).....	258
Xpovio (40 MG Once Weekly).....	258
Xpovio (40 MG Twice Weekly).....	258
Xpovio (60 MG Once Weekly).....	258
Xpovio (60 MG Twice Weekly).....	258
Xpovio (80 MG Once Weekly).....	258
Xpovio (80 MG Twice Weekly).....	258
Xtandi.....	259
Xyrem	260
Y	
Yonsa	261
Z	
Zarxio.....	41, 42

Zejula	262	Ztalmy	266
Zelboraf.....	263	Zurzuvae	267
Zolinza	264	Zydelig	268
Zonisade.....	265	Zykadia Oral Tablet	269

Multi-Language Insert

Multi-language Interpreter Services

English: We have free interpreter services to answer any questions you may have about our health or drug plan. To get an interpreter, just call us at 1- 855-592-7737 (TTY: 711). Someone who speaks English/Language can help you. This is a free service.

Spanish: Tenemos servicios de intérprete sin costo alguno para responder cualquier pregunta que pueda tener sobre nuestro plan de salud o medicamentos. Para hablar con un intérprete, por favor llame al 1-855-592-7737 (TTY: 711). Alguien que hable español le podrá ayudar. Este es un servicio gratuito.

Navajo (Diné): Díí ats'íís dóó azee' bínda'í díłkidgo, Dinék'ehjí yadałti'iigi ła' bich'í' hadíídzih. Béésh bee hane'é t'áá jíík'e be' hódíílnih, 1-855-592-7737 (TTY: 711).

Chinese Mandarin: 我们提供免费的翻译服务，帮助您解答关于健康或药物保险的任何疑问。如果您需要此翻译服务，请致电 1-855-592-7737 (TTY: 711)。我们的中文工作人员很乐意帮助您。这是一项免费服务。

Chinese Cantonese: 您對我們的健康或藥物保險可能存有疑問，為此我們提供免費的翻譯服務。如需翻譯服務，請致電 1-855-592-7737 (TTY: 711)。我們講中文的人員將樂意為您提供幫助。這是一項免費服務。

Tagalog: Mayroon kaming libreng serbisyo sa pagsasaling-wika upang masagotang anumang mga katanungan ninyo hinggil sa aming planong pangkalusugano panggamot. Upang makakuha ng tagasaling-wika, tawagan lamang kami sa 1-855-592-7737 (TTY: 711). Maaari kayong tulungan ng isang nakakapagsalita ng Tagalog. Ito ay libreng serbisyo.

French: Nous proposons des services gratuits d'interprétation pour répondre à toutes vos questions relatives à notre régime de santé ou d'assurance-médicaments. Pour accéder au service d'interprétation, il vous suffit de nous appeler au 1-855-592-7737 (TTY: 711). Un interlocuteur parlant Français pourra vous aider. Ce service est gratuit.

Vietnamese: Chúng tôi có dịch vụ thông dịch miễn phí để trả lời các câu hỏi về chương sức khỏe và chương trình thuốc men. Nếu quý vị cần thông dịch viên xin gọi 1-855-592-7737 (TTY: 711) sẽ có nhân viên nói tiếng Việt giúp đỡ quý vị. Đây là dịch vụ miễn phí.

German: Unser kostenloser Dolmetscherservice beantwortet Ihren Fragen zu unserem Gesundheits- und Arzneimittelplan. Unsere Dolmetscher erreichen Sie unter 1-855-592-7737 (TTY: 711). Man wird Ihnen dort auf Deutsch weiterhelfen. Dieser Service ist kostenlos.

Korean: 당사는 의료 보험 또는 약품 보험에 관한 질문에 대해 드리고자 무료 통역 서비스를 제공하고 있습니다. 통역 서비스를 이용하려면 전화 1-855-592-7737 (TTY: 711). 번으로 문의해 주십시오. 한국어를 하는 담당자가 도와 드릴 것입니다. 이 서비스는 무료로 운영됩니다.

Russian: Если у вас возникнут вопросы относительно страхового или медикаментного плана, вы можете воспользоваться нашими бесплатными услугами переводчиков. Чтобы воспользоваться услугами переводчика, позвоните нам по телефону 1-855-592-7737 (TTY: 711). Вам окажет помощь сотрудник, который говорит по-русски. Данная услуга бесплатная.

Arabic: إننا نقدم خدمات المترجم الفوري المجانية للإجابة عن أي أسئلة تتعلق بالصحة أو جدول الأدوية لدينا. للحصول على مترجم فوري، ليس عليك سوى الاتصال بنا على 1-855-592-7737 (TTY: 711). سيقوم شخص ما يتحدث العربية بمساعدتك. هذه خدمة مجانية.

Hindi: हमारा 4ाTV याT दवाT का या जनाT क बारा मँ आपका ककसा भा ँ क जवाTब दना क िाTाTए हमारा पाTसय. दा, भािाकयाT सा वाए उिापाह. एक दा, भाTिाकयाT ँाT करना क िाकए, बस हमँ 1-855-592-7737 (TTY: 711) पर फा न करँ . का ई D्िाT4 जा कहला ब िााTाT हआपका मदद कर सकताT हँ . यह एक मा, 4 सा वाT हँ .

Italian: È disponibile un servizio di interpretariato gratuito per rispondere a eventuali domande sul nostro piano sanitario e farmaceutico. Per un interprete, contattare il numero 1-855-592-7737 (TTY: 711). Un nostro incaricato che parla Italianovi fornirà l'assistenza necessaria. È un servizio gratuito.

Portuguese: Dispomos de serviços de interpretação gratuitos para responder a qualquer questão que tenha acerca do nosso plano de saúde ou de medicação. Para obter um intérprete, contacte-nos através do número 1-855-592-7737 (TTY: 711). Irá encontrar alguém que fale o idioma Português para o ajudar. Este serviço é gratuito.

French Creole: Nou genyen sèvis entèprèt gratis pou reponn tout kesyon ou ta genyen konsènan plan medikal oswa dwòg nou an. Pou jwenn yon entèprèt, jisrele nou nan 1-855-592-7737 (TTY: 711). Yon moun ki pale Kreyòl kapab ede w. Sa a se yon sèvis ki gratis.

Polish: Umożliwiamy bezpłatne skorzystanie z usług tłumacza ustnego, który pomoże w uzyskaniu odpowiedzi na temat planu zdrowotnego lub dawkowania leków. Aby skorzystać z pomocy tłumacza znającego język polski, należy zadzwonić pod numer 1-855-592-7737 (TTY: 711). Ta usługa jest bezpłatna.

Japanese: 当社の健康 健康保険と薬品 処方薬プランに関するご質問にお答えするために、無料の通訳サービスがあります。通訳をご用命になるには、1-855-592-7737 (TTY: 711). にお電話ください。日本語を話す人者が支援いたします。これは無料のサービスです。